

CARBON DIOXIDE EMISSIONS FROM FROND DECOMPOSITION IN OIL PALM PLANTATIONS ON TROPICAL PEAT

NUR WAKHID^{1*} and TAKASHI HIRANO²

ABSTRACT

Up to now, many studies have been trying to reveal the impact of oil palm plantations on tropical peat on the global carbon (C) balance. Although there are some publications on soil respiration in oil palm plantations on peat soil, information on carbon dioxide (CO₂) emissions from palm litter (pruned fronds) decomposition is limited. Therefore, we quantified the CO₂ emissions through frond decomposition in two different mature oil palm plantations established on peat, an industrial plantation in Riau and a smallholder plantation in Jambi, Indonesia. Frond decomposition was measured using a litter bag method and the decomposition rate constant was determined using a negative exponential equation. Annual CO₂ emissions from frond decomposition were estimated at 1.48 and 1.12 Mg C ha⁻¹ yr⁻¹, respectively, in industrial and smallholder plantations. As a result, CO₂ emissions from frond decomposition accounted for 10%-14% of heterotrophic respiration from mature oil palm plantations on tropical peat.

Keywords: industrial plantation, mature plantation, palm litter, smallholder plantation.

Received: 17 April 2022; **Accepted:** 21 September 2022; **Published online:** 19 October 2022.

INTRODUCTION

Tropical peatland has become one of the most efficient carbon sinks on the earth and plays an important role in the global carbon (C) cycle. This ecosystem mostly concentrated in Southeast Asia (SEA), the upper Amazon and in Congo Basin, covers only 16% of global peatland areas but is stored up to 105 Pg of C (Dargie *et al.*, 2017; Page *et al.*, 2011). However, this huge C deposit is now becoming vulnerable to being changed as a massive C source due to the land conversion that promotes peat decomposition and peat fires (Hooijer *et al.*, 2012; Miettinen *et al.*, 2017).

Driven by population growth and economic development, during the last two decades, many areas of tropical peat swamp forest in SEA have been converted to agricultural plantations, mostly for oil palm and acacia trees (Miettinen *et al.*, 2012). The total area of peat swamp forest in Peninsular Malaysia, Sumatra and Borneo has been estimated to decrease from 64 x 10⁵ ha in 2007 to 46 x 10⁵ ha, while oil palm plantations on peatland had extended up to 31 x 10⁵ ha in 2015 (Miettinen *et al.*, 2016). The global demand for palm oil in the world is expected to double by 2030 compared to 2000 (Yan, 2017). Thus, the impact of this plantation on the global C balance could be large.

To date, there are several publications on oxidative peat decomposition or heterotrophic respiration in oil palm plantations on peat (*e.g.* Dariah *et al.*, 2014; Hergoualc'h *et al.*, 2017; Ishikura *et al.*, 2018; Manning *et al.*, 2019; Melling *et al.*, 2013 and Wakhid and Hirano, 2021a). Unfortunately, none of these have studied carbon dioxide (CO₂)

¹ Research Center for Ecology and Ethnobiology, National Research and Innovation Agency, 16911, Cibinong, Bogor, Indonesia.

² Research Faculty of Agriculture, Hokkaido University, 060-8589, Sapporo, Japan.

* Corresponding author e-mail: n_wakhid@yahoo.com

emissions through palm litter (pruned fronds) decomposition, except Wakhid and Hirano (2021a), which conducted the study on a young oil palm plantation (7 years old). More field experiments are necessary, particularly in mature oil palm plantations to understand the contribution of CO₂ emissions through frond decomposition to the C balance of oil palm plantations on peat. The objective of this study was to quantify how much CO₂ is emitted from the decomposition of pruned fronds left on the ground in mature oil palm plantations on peat soil.

MATERIALS AND METHODS

The study was conducted over one year from February 2018 to March 2019, in mature oil palm (*Elaeis guineensis* Jacq.) plantations in Sumatra, Indonesia. One site was an industrial plantation in Riau (0° 42' 16" N, 101° 42' 52.8" E), and another site was a smallholder plantation in Jambi (1° 14' 20" S, 103° 35' 23" E). The oil palm cultivar in Riau was Marihat and in Jambi was mixed between the Marihat and Socfin with a ratio of 4:1. Thus, weighting factors of 0.8 and 0.2 were applied to the data for Marihat and Socfin, respectively, for the calculation of the CO₂ emissions in Jambi. Further information about the study sites was described in Wakhid *et al.* (2022).

To estimate the decomposition, fresh fronds were cut down, separated into two parts (tip and base), and put into litter bags. The litter bags were 40 × 80 cm, with a mesh size of 2 mm. Before being inserted into the litter bags, the fresh weight of each frond was recorded. Each bag contained about 800 g of frond. Also, the initial values of dry weight and C content of fronds were measured (n = 3). In Riau, the litter bags were set on the top of frond heaps in February 2018 with a total of 100 bags for each tip and base. The bags were retrieved in September 2018, January 2019 and June 2019, but only 99 bags were in total because most of them were missing. In Jambi, 36 bags of tips and bases for each of Marihat and Socfin were set equally on the top of and under frond heaps, in February 2018. In total, six bags of each tip and base from the two positions were retrieved in September 2018, February 2019 and July 2019. Each litter bag was cleaned after being retrieved, and the dry weight and C content were measured.

The dry weight was measured by oven drying for 48 hr at 70°C. The C content was estimated by the loss of ignition method, using a conversion factor from organic matter to organic C of 0.58 (Agus *et al.*, 2011). The decomposition rate constant (*k*) was estimated using a negative exponential equation from the plot of the residual ratio of C amount (*Y*) against elapsed time (*t*) following Moradi *et al.* (2014) as shown in Equation (1).

$$Y = Y_0 \exp(-kt) \quad (1)$$

where C loss through frond decomposition was estimated using the *k* and initial C amount (*Y*₀). Then, the annual CO₂ emission was calculated using pruned frond production × monthly amount of CO₂ emissions. The annual C input was estimated from annual pruned fronds × tree density × dry weight × C content. Statistical analysis (ANOVA, Tukey HSD test, and exponential graph) were conducted using Excel and R software (R Development Core Team 2019, version 3.5.3). Further detail about CO₂ emissions calculation was described in Wakhid and Hirano (2021a; 2021b).

RESULTS AND DISCUSSION

The C content of pruned fronds in Riau and Jambi was in the range of 53%-57% (Table 1). C and Nitrogen (N) contents of leaf were significantly different with rachis (Table 1). However, the C and N contents of the tip and base parts were not significantly different. CN ratio was significantly higher in rachis than in leaf. Socfin in Jambi has the lowest CN ratio than that of Marihat in Riau and Jambi, respectively (Table 1).

In Riau, the amount of litter bags in the second and third retrieving was smaller than that of the first retrieving (n = 45, 27 and 27 litter bags, respectively, for the first, second and third retrieving) because several bags were lost during the study. Thus, the *k* value was calculated using all retrieved samples without tip and base parts separation to be 1.86 yr⁻¹ (Figure 1). The annual C input was 2.54 Mg C ha⁻¹ yr⁻¹, and CO₂ emissions from frond decomposition were estimated at 1.48 Mg C ha⁻¹ yr⁻¹ (Table 2).

In Jambi, although, the *k* values among tip and base parts and between two cultivars were not significantly different, CO₂ emissions through frond decomposition should be different among the cultivars because the size of Socfin was 1.4 times larger than that of Marihat (Table 1). Thus, the *k* values were calculated using all data for each cultivar (n = 12), to be 1.57 and 1.81 yr⁻¹, respectively, for Marihat and Socfin (Figure 1). The annual C input and CO₂ emissions from pruned fronds were estimated at 2.07 and 1.12 Mg C ha⁻¹ yr⁻¹, respectively (Table 2).

CO₂ emissions from frond decomposition in this study were higher than in a young oil palm plantation in Banjarbaru (Wakhid and Hirano (2021a): 0.38 Mg C ha⁻¹ yr⁻¹). This discrepancy was most likely caused by the production of frond that was larger in mature than in young plantations. The CO₂ emissions (1.48 and 1.12 Mg C ha⁻¹ yr⁻¹, in Riau and Jambi, respectively) accounted for 14% and 11% of heterotrophic respiration from mature oil palm

TABLE 1. INITIAL AMOUNT OF CARBON (C), NITROGEN (N), CN RATIO, AND WATER CONTENT (WC) OF FROND (MEAN ± 1 SD, n = 3).

Samples	C (%)	N (%)	CN ratio	WC (%)
Marihat (M) in Riau (dry weight = 2578 ± 509 g frond ⁻¹ , n = 5)				
Leaf (tip)	54.60 ± 0.54a	1.23 ± 0.58a	50.30 ± 19.10a	59.30 ± 0.30
Rachis (tip)	56.70 ± 0.28b	0.38 ± 0.13b	160.90 ± 58.90b	60.20 ± 8.30
Leaf (base)	55.00 ± 0.29a	1.62 ± 0.44a	35.40 ± 8.40a	59.10 ± 1.60
Rachis (base)	55.60 ± 0.35b	0.39 ± 0.06b	143.70 ± 20.00b	37.80 ± 6.60
Marihat (M) in Jambi (dry weight = 2254 ± 188 g frond ⁻¹ , n = 5)				
Leaf (tip)	54.00 ± 0.33a	1.90 ± 0.33a	29.20 ± 5.70a	54.90 ± 2.70
Rachis (tip)	57.10 ± 0.07b	0.66 ± 0.08b	87.20 ± 11.40b	65.90 ± 4.20
Leaf (base)	54.40 ± 0.40a	2.00 ± 0.19a	27.40 ± 2.30a	58.50 ± 2.30
Rachis (base)	56.90 ± 0.25b	0.60 ± 0.10b	96.20 ± 15.40b	71.10 ± 7.50
Socfin (S) in Jambi (dry weight = 3387 ± 567 g frond ⁻¹ , n = 5)				
Leaf (tip)	53.10 ± 0.89a	2.27 ± 0.16a	23.50 ± 1.50a	50.10 ± 11.40
Rachis (tip)	56.80 ± 0.09b	0.75 ± 0.13b	77.00 ± 13.30b	69.80 ± 3.70
Leaf (base)	53.60 ± 1.32a	2.15 ± 0.51a	25.90 ± 6.00a	56.50 ± 2.60
Rachis (base)	56.40 ± 0.11b	0.79 ± 0.20b	74.90 ± 20.60b	71.10 ± 4.80
ANOVA (<i>p</i> - value)				
Component (leaf vs. rachis)	<0.001	<0.001	<0.001	0.210
Position (tip vs. base)	0.76	0.64	0.68	0.50
Site (<i>M</i> in Riau vs. <i>M</i> in Jambi)	0.74	0.18	0.08	0.13
Cultivar (<i>M</i> vs. <i>S</i> in Jambi)	0.42	0.06	<0.05	0.16
Interaction (component-position)	<0.05	0.62	0.98	0.06

Note: Data of C (%) of Marihat and Socfin in Jambi was adapted from Wakhid and Hirano (2021b). Different letters denote significant differences between leaf and rachis.

TABLE 2. CARBON (C) INPUT AND CARBON DIOXIDE (CO₂) EMISSIONS FROM FROND DECOMPOSITION IN RIAU AND JAMBI

Site and cultivar	Tree density (trees ha ⁻¹)	Frond production (dry weight, Mg ha ⁻¹ yr ⁻¹)	C input (Mg C ha ⁻¹ yr ⁻¹)	CO ₂ emissions (Mg C ha ⁻¹ yr ⁻¹)
Industrial in Riau (<i>Marihat</i>)	148	4.58	2.54	1.48
Smallholder in Jambi	125	3.72	2.07	1.12
<i>Marihat</i>	100	2.71	1.51	0.80
<i>Socfin</i>	25	1.02	0.56	0.32




Figure 1. C loss pattern of a) *Marihat* in Riau and b) *Marihat* (squares, dash line) and *Socfin* (circles, solid line) in Jambi. An exponential curve is fitted for each retrieval time. Riau: $Y = e^{-1.86x}$ ($r^2 = 0.83$), and Jambi: $Y_m = e^{-1.57x}$ ($r^2 = 0.90$), $Y_s = e^{-1.81x}$ ($r^2 = 0.84$). Y_m and Y_s represent the equations for *Marihat* and *Socfin* cultivars, respectively.

plantations on tropical peat (Dariah *et al.* (2014): 15 years old, 10.4 Mg C ha⁻¹ yr⁻¹). Also, accounted for 13% and 10 %, in Riau and Jambi, respectively, to a default CO₂ emission factor Tier 1 methodology of oil palm plantation on peat soil (11 Mg C ha⁻¹ yr⁻¹) from the Intergovernmental Panel on Climate Change (IPCC, 2014). Following the amount of CO₂ emissions, the contribution in mature was larger than in young oil palm plantations (Wakhid and Hirano (2021a): Accounting for 2.4% of heterotrophic respiration).

Some data in Jambi from February 2018 to February 2019 was presented at International Conference on Sustainable Tropical Land Management (Wakhid and Hirano, 2021b). Unexpectedly, CO₂ emission in Jambi from February 2018 to February 2019 (Wakhid and Hirano (2021b): 1.09 Mg C ha⁻¹ yr⁻¹) was similar to February 2018 to September 2019 (this study: 1.12 Mg C ha⁻¹ yr⁻¹). Frond decomposition could likely be estimated in a short study as long as the disturbance and environmental changes are minimal. To our knowledge, previous studies on CO₂ emissions from frond decomposition were limited, thus more studies for comparison were not found.

The annual C input from pruned fronds and *k* values from this study were compared with some previous studies. Annual C input in this study (2.54 and 2.07 Mg C ha⁻¹ yr⁻¹, respectively, in Riau and Jambi) was lower than the C input of pruned fronds from oil palm on mineral soil in Jambi (Kotowska *et al.* (2016): 2.89 Mg C ha⁻¹ yr⁻¹). However, our result was higher than that of 1.33-1.79 Mg C ha⁻¹ yr⁻¹ of C input from pruned fronds in an oil palm plantation on mineral soil in North Sumatra (Lamade and Bouillet, 2005). *k* values of Marihat in Riau and Socfin in Jambi (Figure 1) were similar to the *k* value of pruned fronds from oil palms on mineral soil in Malaysia (1.80 yr⁻¹; Moradi *et al.*, 2014), but higher than that of another study also in Malaysia (0.73 yr⁻¹; Khalid *et al.*, 2000).

CONCLUSION

CO₂ emissions from frond decomposition were investigated in mature oil palm plantations on tropical peat, industrial and smallholder plantations, in Riau and Jambi. Using decomposition rate constant (*k*), initial carbon (C) amount, and pruned frond production, CO₂ emissions from frond decomposition were estimated to be 1.48 and 1.12 Mg C ha⁻¹ yr⁻¹, respectively, in Riau and Jambi. These CO₂ emissions accounted for 10%-14% of heterotrophic respiration in oil palm plantations on peat. The contribution of frond decomposition is not large but should be considered in peat soil respiration calculation.

ACKNOWLEDGEMENT

We thank Fahmuddin Agus and Ai Dariah for permitting us to use their field research for the study. This study was supported by Indonesian Agency for Agricultural Research and Development and JSPS KAKENHI (No.17H01477, 18H02238, and 19H05666).

REFERENCES

- Agus, F; Hairiah, K and Mulyani, A (2011). *Measuring Carbon Stock in Peat Soils: Practical Guidelines*. World Agroforestry Centre (ICRAF) Southeast Asia Regional Program, Indonesian Centre for Agricultural Land Resources Research and Development. Bogor, Indonesia. 78 pp.
- Dargie, G C; Lewis, S L; Lawson, I T; Mitchard, E T A; Page, S E; Bocko, Y E and Ifo, S A (2017). Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature*, 542: 86-90. DOI: 10.1038/nature21048.
- Dariah, A; Marwanto, S and Agus, F (2014). Root- and peat-based CO₂ emissions from oil palm plantations. *Mitig. Adapt. Strateg. Glob. Chang.*, 19: 831-843. DOI: 10.1007/s11027-013-9515-6.
- Hergoualc'h, K; Hendry, D T; Murdiyarso, D and Verchot, L V; (2017). Total and heterotrophic soil respiration in a swamp forest and oil palm plantations on peat in Central Kalimantan, Indonesia. *Biogeochemistry*, 135: 203-220. DOI: 10.1007/s10533-017-0363-4.
- Hooijer, A; Page, S; Jauhiainen, J; Lee, W A; Lu, X X; Idris, A and Anshari, G (2012). Subsidence and carbon loss in drained tropical peatlands. *Biogeosciences*, 9(3): 1053-1071. DOI: 10.5194/bg-9-1053-2012.
- IPCC (2014). *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories*. IPCC, Wetlands, Switzerland. 354 pp.
- Ishikura, K; Hirano, T; Okimoto, Y; Hirata, R; Kiew, F; Melling, L; Aeries, E B; Lo, K S; Musin, K K; Waili, J W; Wong, G X and Ishii, Y (2018). Soil carbon dioxide emissions due to oxidative peat decomposition in an oil palm plantation on tropical peat. *Agric. Ecosyst. Environ.*, 254: 202-212. DOI: 10.1016/j.agee.2017.11.025.
- Khalid, H; Zin, Z Z and Anderson, J M (2000). Decomposition processes and nutrient release patterns of oil palm residues. *J. Oil Palm Res.*, 12(1): 46-63.

- Kotowska, M M; Leuschner, C; Triadiati, T and Hertel, D (2016). Conversion of tropical lowland forest reduces nutrient return through litterfall and alters nutrient use efficiency and seasonality of net primary production. *Oecologia*, 180: 601-618. DOI: 10.1007/s00442-015-3481-5.
- Lamade, E and Bouillet, J P (2005). Carbon storage and global change: The role of oil palm. *Oléagineux Corps Gras Lipides (OCL)*, 12: 154-160. DOI: 10.1051/ocl.2005.0154.
- Manning, F C; Kho, L K; Hill, T C; Cornulier, T and Teh, Y A (2019). Carbon emissions from oil palm plantations on peat soil. *Front. For. Glob. Change*, 2: 37, 21. DOI: 10.3389/ffgc.2019.00037.
- Melling, L; Yun Tan, C S; Goh, K J and Hatano, R (2013). Soil microbial and root respirations from three ecosystems in tropical peatland of Sarawak, Malaysia. *J. Oil Palm Res.*, 25: 44-57.
- Miettinen, J; Hooijer, A; Shi, C; Tollenaar, D; Vernimmen, R; Liew, S C; Malins, C and Page, S E (2012). Extent of industrial plantations on Southeast Asian peatlands in 2010 with analysis of historical expansion and future projections. *Glob. Change Biol. Bioenergy*, 4: 908-918. DOI: 10.1111/j.1757-1707.2012.01172.x.
- Miettinen, J; Shi, C and Liew, S C (2016). Land cover distribution in the peatlands of Peninsular Malaysia, Sumatra and Borneo in 2015 with changes since 1990. *Glob. Ecol. Conserv.*, 6: 67-78. DOI: 10.1016/j.gecco.2016.02.004.
- Miettinen, J; Hooijer, A; Vernimmen, R; Liew, S C and Page, S E (2017). From carbon sink to carbon source: Extensive peat oxidation in insular Southeast Asia since 1990. *Environ. Res. Lett.*, 12: 024014.
- Moradi, A; Teh, C B S; Goh, K J; Husni, M H A and Ishak, C F (2014). Decomposition and nutrient release temporal pattern of oil palm residues. *Ann. Appl. Biol.*, 164(2): 208-219. DOI: 10.1111/aab.12094.
- Page, S E; Rieley, J O and Banks, C J (2011). Global and regional importance of the tropical peatland carbon pool. *Glob. Change Biol.*, 17: 798-818. DOI: 10.1111/j.1365-2486.2010.02279.x.
- Wakhid, N and Hirano, T (2021a). Soil CO₂ emissions and net primary production of an oil palm plantation established on tropical peat. *Mires Peat*, 27: 11. DOI: 10.19189/MaP.2021.OMB.StA.2159.
- Wakhid, N and Hirano, T (2021b). Contribution of CO₂ emission from litter decomposition in an oil palm plantation on tropical peatland. *1st International Conference on Sustainable Tropical Land Management. IOP Conf. Series: Earth and Environmental Science*. 648: 012133. DOI: 10.1088/1755-1315/648/1/012133.
- Wakhid, N; Hirano, T; Dariah, A and Fahmuddin, A (2022). Net primary production of oil palm plantations on tropical peat. *Mires and Peat*, 28: 02-12. DOI: 10.19189/MaP.2021.SNPG.StA.2288.
- Yan, W (2017). A makeover for the world's most hated crop. *Nature*, 543(16): 306-308.