

STUDY ON THE EFFECT OF VARIOUS TYPES OF FERTILISER ON THE PRODUCTION OF OIL PALM ROOT CUTTING

CECEP IJANG WAHYUDIN¹; HARIYADI^{1*}; SUDRAJAT¹; SUDIRMAN YAHYA¹ and SYAIFUL ANWAR²

ABSTRACT

In the context of Indonesia, the cultivation of oil palm (*Elaeis guineensis*) is extremely significant and the application of diverse fertiliser variants can affect crop yield through the implementation of root-cutting methods. The modification of root morphology is also a strategy to address the issue of suboptimal nutrient uptake and soil physical properties exert a significant influence on the development. Therefore, this study used a nested plot design as the main plot, comprising plants aged 7, 12, and 16 years. The subplots were demarcated by fertiliser types, including control (no fertilisation), single fertiliser K (2.25 kg KCl/tree), palm frond litter (65 kg/tree), and empty fruit bunches (65 kg/tree). Furthermore, the treatment was replicated 4 times, resulting in a total of 48 experimental units. The results showed that plants aged 7 years exhibited the most robust growth compared to 12 and 16 years, across all fertiliser types, as indicated by stem circumference, stem diameter, and NDVI values. Regarding the primary, secondary, and tertiary optimal root distribution, 7 year old plants responded to K fertiliser. However, tertiary distribution did not significantly differ from 12 year-old plants when K fertiliser was applied. The nutrient levels in 7 year old and 12 year old plants varied in terms of pH, organic carbon (C-organic), and potassium (K).

Keywords: fertility soil, oil palm, root cutting.

Received: 20 July 2023; **Accepted:** 29 December 2023; **Published online:** 4 April 2024.

INTRODUCTION

Oil palm (*Elaeis guineensis* Jacq.) is a significant plantation crop in the agricultural sector. Among different oil-producing plants, it holds economic value as a foreign exchange source and Indonesia, Malaysia, and Thailand are the world's largest producers. Several activities are conducted to enhance palm oil production, including expansion, rehabilitation of existing plantations, and intensification (Nurchahyo and Ariani, 2017). Fertilisation can be applied to oil palm plants subjected to root pruning to improve their production and quality.

Root pruning can result in drought stress for plants due to disrupted nutrient and water absorption. It can also inhibit stem and shoot diameter growth (Fini *et al.*, 2013; Watson, 1998). However, root pruning enhances plant nutrient uptake through morphological and physiological mechanisms. Morphological mechanisms include root hair distribution and formation, where cutting roots stimulate the growth of new ones at the cut ends (Miller and Neely, 1993). Root pruning increases the percentage of dry-weight growth and root dry weight in oil palms (Luok and Raharjo, 2017). Physiological mechanisms for improving nutrient uptake through root include root kinetics and nutrient mobilisation through root exudation (Rao *et al.*, 1999). Root pruning is performed on different plants with varying distances from the stem, pruning depth, and intensity. In the case of oil palm, it possesses a fibrous root system, with tertiary and quaternary roots playing a crucial role in nutrient and water absorption (Yahya *et al.*, 2010). This condition

¹ Department of Agronomy and Horticulture, IPB University, Bogor, 16680 Indonesia.

² Department of Soil Science and Land Resources, Faculty of Agriculture, IPB University, Bogor, 16680 Indonesia.

* Corresponding author e-mail: hariyadibdp@apps.ipb.ac.id

leads to changes in organic acids expressed by roots as plants adapt to enhance nutrient uptake and optimise the absorption of essential nutrients when provided with the right types of fertiliser.

The application of fertiliser can enhance the production and quality of root-pruned plants. Proper application provides the necessary nutrients for optimal growth and development of plants. Previous studies have shown that providing fertiliser with the appropriate concentration, composition, and dosage can increase plant productivity. One nutrient that aids in plant recovery after drought stress is potassium, enhancing plant resilience to drought stress. This is influenced by physiological and biochemical processes related to the synthesis of compounds such as proline, antioxidant activity, and phenols. Under drought stress conditions, the osmotic pressure in the environment is higher than inside plant cells, making it difficult for plants to absorb water. Plants increase the concentration of intracellular solutes such as proline and reduce the intracellular osmotic potential to maintain potential water balance (Cha-um *et al.*, 2010). These compounds actively contribute to the physiological mechanisms that enable plants to withstand drought stress (Egilla *et al.*, 2001). Additionally, using organic and potassium (K) fertiliser offers more environmentally friendly options providing additional benefits to root-pruned plants. Organic fertiliser derived from well-packaged oil palm plant waste can serve as a nutrient source.

The waste generated by oil palm plantations can manifest as solid waste, including empty fruit bunches (EFB), palm fronds, and palm leaves. Sarwono (2008) states the advantages of EFB compost, such as the high K content, absence of starter and chemicals, enrichment of existing soil nutrients, and improvement of physical, chemical, and biological properties. This can be used as organic mulch or incorporated into soil to enhance soil fertility. The physical properties of soil are one of the factors that influence root growth, and inadequate physical properties can impede root development (Andiyarto and Purnomo, 2012). Habi (2015) explains that the increase in soil pore space occurs due to the stimulation of soil aggregates by organic matter, resulting in a decrease in soil bulk density. Additionally, Widodo and Kusumo (2018) state that the increase in organic matter content acts as a binder in soil aggregate formation, leading to the creation of more inter-aggregate (macropores) and intra-aggregate spaces (micropores), enhancing air and water availability in soil.

Applying organic matter increases the total pore space, reduces soil volume (Greenland, 2003), and improves soil porosity by 13% (Endriani *et al.*, 2003). Oil palm plantations possess abundant organic matter in the form of leaf litter and empty

fruit bunches. Oil palm EFB is an abundant waste product derived from palm oil mills. This is known to improve the physical, chemical, and biological properties of soil, in addition to having a high nutrient content. Sarwono (2008) reports that EFB compost contains essential plant nutrients, including 1.5% nitrogen (N), 0.5% phosphorus (P), 7.3% potassium (K) and 0.9% magnesium (mg). Furthermore, Hayat and Andayani (2014) indicate that EFB compost's nutrient content includes total N (1.91%), K (1.51%), calcium (Ca) (0.83%), P (0.54%), Mg (0.09%), organic carbon (51.23%), C/N ratio (26.82%) and a pH of 7.

EFB compost serves a dual function by adding nutrients to soil and increasing its organic matter content crucial for improving the physical properties of soil. The increased organic matter enhances the stability of soil structure and improves its water retention capacity. These improvements in soil's physical properties positively impact root growth and nutrient uptake by plants (Rozy *et al.*, 2013). Apart from EFBs, there is a significant accumulation of palm frond litter around oil palm trees. Therefore, palm frond litter has the potential to serve as a source of organic matter to improve the physical and chemical properties of soil in oil palm plantations. Using palm frond litter as fertiliser for oil palm trees is viable since it contains an average nutrient content of 1.67% N, 0.17% P, 0.23% K, 0.83% Ca and 0.26% Mg (Yuniati and Widyastuti, 2014).

The application of palm frond litter and EFBs aims to increase the organic matter content, improving the physical properties of soil. The use of empty fruit bunches is a common practice in oil palm plantations. However, the comprehensive exploration of palm frond litter, particularly in the context of its impact on root pruning, remains an area that has not received extensive attention. EFB compost, derived from palm oil mills, is a readily available waste product known to enhance soil's physical, chemical, and biological properties. The compost is abundant throughout the year, accounting for approximately 20%-27% of processed fresh fruit bunches (FFB) (Baryanto *et al.*, 2015). Increasing the dosage has been found to correlate with an increase in root volume (Amri *et al.*, 2018). The advantages of compost derived from palm frond litter include its high K content, the use of bio decomposers obtained from microbial isolates found beneath palm frond stacks in the field, the enrichment of soil nutrients, and the ability to improve the physical, chemical, and biological properties of soil (Yuniati and Widyastuti, 2014).

Using palm waste as fertiliser holds great potential for enhancing the productivity of root-pruned crops. Therefore, this study aims to enhance the understanding of how root-pruned crops respond to various fertilisers. It provides valuable information for farmers and agricultural

practitioners to improve production efficiency and promote agricultural sustainability in the context of root-pruned crops. By understanding the nutrient requirements of these crops and applying suitable fertiliser, farmers can enhance the quality and quantity of their root-pruned crop yields.

MATERIALS AND METHODS

The study was conducted from October 2022 to May 2023 in Teluk Merbau Village, Dayun District, Siak Sriindrapura Regency, Riau. Soil nutrient analysis was performed at the Department of Soil Science and Land Resources, Faculty of Agriculture, IPB University. The study used K fertiliser, palm frond litter, and empty fruit bunches as materials. Furthermore, the tools employed included a measuring tape, hoe, soil auger, digital scale, SPAD-502, Haga Altimeter, and GPS. The study followed a nested design, with plant age as the main plot and fertiliser types as the subplot. The plant age factor consisted of three levels, namely 7, 12 and 16 years, while the fertiliser types factor had four levels, including Control (without fertilisation), single K fertiliser (2.25 kg KCl/tree), palm frond litter (65 kg/tree) and EFBs (65 kg/tree). Each treatment was replicated four times, resulting in 48 experimental units of four plants, totalling 192 (Darmosarkoro *et al.*, 2008).

The data were analysed using analysis of variance (ANOVA) with SAS software at a 95% confidence level ($\alpha = 5\%$). Duncan's Multiple Range Test (DMRT) was conducted at a 95% confidence level ($\alpha = 5\%$) when analysis of variance indicated a significant treatment effect. The observed variables included plant height, measured from the ground surface to the rudimentary spine on the front using a Haga Altimeter. Stem circumference and diameter were measured with a tape by

calculating the average circumference at the base of the stem. Root distribution was assessed after pruning, and samples were taken from one plant in each experimental unit, resulting in 28 sampling points. The number and weight of fruit bunches were observed at 1, 2, 3, 4, 5, and 6 months after the treatment. Furthermore, soil nutrient observations and natrium with Kjeldhal method, phosphor with bray I, and kalium with washing method using ammonium acetate (NH₄OAc) at pH 7 in root were conducted.

RESULTS AND DISCUSSION

Agronomic and Production Characteristics

The analysis of variance showed a significant variation in the height of oil palm plants due to the application of different fertilisers at the beginning and end of the observation period for each plant age, as shown in *Table 1*.

Table 1 shows that the tallest plant height was observed in 16 year old plants compared to those aged 12 and 7. Among the different types of fertiliser used, including the control, K fertiliser, midrib litter, and empty fruit bunches, there were no significant differences at each plant age. Generally, the height of oil palm plants increased from the initial to the final for each combination of plant age and fertiliser types. A higher plant height indicates better growth; as plants age, their height tends to increase compared to younger plants.

The application of different fertilisers did not significantly impact plant height, resulting in no differences observed among the treatments. Several factors influence the availability of nutrients in the soil for plant absorption and use. Fangeria *et al.* (2009) state that climate, soil conditions, plant-related factors, and interactions influence nutrient

TABLE 1. OIL PALM PLANT HEIGHT FOR DIFFERENT PLANTING AGES AND THE APPLICATION OF VARIOUS FERTILISER

Plant age (yr)	Types of fertiliser	Initial plant height		Final plant height	
7	Control	343.06	c	373.06	c
	K fertiliser	352.56	c	382.56	c
	Leaf litter	350.69	c	380.69	c
	Empty bunches	338.06	c	368.06	c
12	Control	610.31	b	642.31	b
	K fertiliser	645.50	b	677.50	b
	Leaf litter	629.50	b	661.50	b
	Empty bunches	643.25	b	675.25	b
16	Control	712.31	a	752.31	a
	K fertiliser	747.50	a	787.50	a
	Leaf litter	731.50	a	771.50	a
	Empty bunches	745.25	a	785.25	a

availability. Additionally, Syakir and Gusmaini (2012) emphasise the importance of considering factors that ensure plant nutrient availability for growth and development. Nutrients play a vital role in the formation of plant tissue, and any imbalances in soil nutrient levels can disrupt this process.

Even though fertiliser treatments did not yield significant differences, applying K fertiliser resulted in taller plants compared to other types across different ages of oil palm plants. K fertiliser contains the essential macronutrient, which is crucial for various physiological processes in plants, including protein synthesis, photosynthesis, cell osmotic pressure regulation, and water and sugar balance control. Applying K fertiliser to oil palm can increase growth, such as plant height, productivity, and fruit quality.

Based on the analysis of variance, there were significant differences in oil palm trunk circumference among different plant ages. The largest trunk circumference, both at the beginning and end of the measurement, was observed in plants aged seven years, as shown in *Table 2*.

The trunk circumference of oil palm aged seven years showed a greater circumference than aged 12 and 16 years at the beginning and end of the observations. The use of different types of fertiliser at each age did not show significant differences for control, K fertiliser, midrib leaf litter and empty fruit bunches.

The diameter of the oil palm stem from the analysis of variance reported a significant difference in the diameter of the stem at each age. The largest diameter was shown in aged seven compared to 12 and 16 years old. The application of different types of fertiliser at each age showed no significant difference, as reported in *Table 3*.

The diameter of the stem is an indicator that can be used to determine plant growth and development. Oil palms aged seven years have a larger trunk diameter than others. Plants aged seven years experience faster growth in the vegetative phase using various types of fertiliser, which can spur growth in stem diameter. The application of various types of fertiliser is more optimally used in oil palms aged seven compared to plants aged 12 and 16 years, hence the diameter is more significant.

TABLE 2. OIL PALM TRUNK CIRCUMFERENCE FOR VARIOUS PLANTING AGES AND APPLICATION OF DIFFERENT FERTILISER

Plant age (yr)	Types of fertiliser	Initial trunk circle		Final trunk circle	
7	Control	350.43	a	353.43	a
	K fertiliser	362.50	a	365.50	a
	Leaf litter	360.18	a	363.18	a
	Empty bunches	354.43	a	357.43	a
12	Control	294.43	b	298.43	b
	K fertiliser	306.50	b	310.50	b
	Leaf litter	304.18	b	306.31	b
	Empty bunches	298.43	b	302.43	b
16	Control	259.43	c	263.43	c
	K fertiliser	271.50	c	275.50	c
	Leaf litter	267.31	c	271.31	c
	Empty bunches	263.43	c	267.43	c

TABLE 3. OIL PALM TRUNK DIAMETER FOR VARIOUS TYPES OF PLANTING AGES AND THE APPLICATION OF VARIOUS TYPES OF FERTILISER

Plant age (yr)	Types of fertiliser	Initial bar diameter		Final bar diameter	
7	Control	111.62	a	112.56	a
	K fertiliser	115.50	a	116.50	a
	Leaf litter	113.62	a	114.81	a
	Empty bunches	112.87	a	113.56	a
12	Control	93.81	b	95.00	b
	K fertiliser	97.62	b	98.81	b
	Leaf litter	96.81	b	97.65	b
	Empty bunches	95.06	b	96.37	b
16	Control	82.65	c	83.87	c
	K fertiliser	86.50	c	87.62	c
	Leaf litter	85.06	c	86.56	c
	Empty bunches	83.87	c	85.18	c

Oil palm fruit weight from the analysis of variance showed a significant difference to the observations at the 1st, 2nd, 3rd, 4th, 6th, 7th and 8th months. During the 5th month of observation, there was no significant difference in using different fertilisers at various plant ages, as shown in *Table 4*.

The highest total fruit weight of oil palms was found in oil palm aged seven years with the application of K fertiliser and midrib leaf litter and in plants aged 12 years with the application of empty fruit bunches. These three types of fertiliser contributed essential nutrients, resulting in an augmentation of the weight of oil palm fruit. Kasem *et al.* (2010) stated that using nitrogen fertiliser combined with P and K and organic fertiliser increased oil palm yields and fruit colour, weight, weight and length.

Based on the analysis of variance, the number of oil palm bunches showed no significant difference in the age of the plants and the application of various types of fertiliser, as shown in *Table 5*.

The bunches are the number of fruits produced by oil palms. The number shows no significant difference with applying various types of fertiliser at different ages of oil palms. Appropriate fertiliser application can affect the growth and productivity of oil palms and the number of bunches produced. Fertilisers rich in N, P, and K can be absorbed by plants, influencing the growth and formation of bunches (Syahfitri, 2008).

The formation of oil palm bunches is not solely contingent on fertiliser application, a multitude of factors exert influence, including genetic factors and environmental conditions. Genetic factors, including plant varieties, play an important role in determining the number of bunches produced, with certain varieties exhibiting superior bunch production owing to inherent genetic traits related to bunch formation. Furthermore, environmental conditions, such as the availability of water, sunlight exposure, and nutrient levels, exert a profound influence on the proliferation of oil palm bunches. Plants thriving under optimal conditions, characterised by water resources, sufficient sunlight exposure, and a well-balanced nutrient supply, tend to manifest an enhanced propensity for bunch production.

Analysis of variance showed significant differences in the distribution of oil palm root based on the age and various types of fertiliser applications on the distribution of primary, secondary, and tertiary roots. There were no significant differences in the plant's age and the application of various types of fertiliser, as shown in *Table 6*.

The distribution of roots reported significant differences among primary, secondary, and tertiary roots. *Table 6* shows that the overall distribution

of roots is influenced by the age of the plant and the types of fertiliser used. The use of K fertiliser significantly affected the distribution of roots at various ages of the plants. Fertiliser had the highest primary, secondary, and tertiary root distribution values for oil palms aged seven years, even for plants aged 12 and 16 years. It also affected the distribution of roots of oil palms aged 12 and 16 years but towards secondary and tertiary roots (Syahfitri, 2008).

The use of K fertiliser increased the distribution of primary, secondary, and tertiary roots in oil palms. K fertiliser application also affected root distribution by increasing root growth. K fertiliser stimulated primary root growth, cell division, and extension of primary root, which supported plant growth and development. Another function was to influence plant growth hormones such as auxin, which affected root formation and increased the distribution of roots.

Analysis of variance showed a significant difference in the Normalised Difference Vegetation Index (NDVI) for plants of different ages, with the highest value found in plants aged seven years. However, there was no significant difference for plants aged 12 and 16 years, as shown in *Table 7*.

The NDVI index is a form of vegetation index used to evaluate, and monitor plant conditions and calculate the ratio of the spectral values of the red and near-infrared bands reflected from plants (Cahyono *et al.*, 2019; Yuniasih, 2022). A 7-year-old oil palm with a high NDVI value compared to the age of other plants at 0.736 -0.756 indicates that the plant is in healthy condition and actively carrying out photosynthesis. Furthermore, the plant has a sufficient amount of chlorophyll and can absorb red light optimally for the process. Healthy plants also have dense, healthy cells, affecting near-infrared light's reflection. Oil palm aged seven years shows that the plants have sufficient chlorophyll content and dense cells, reflecting good photosynthetic activity and optimal growth conditions. According to (Foong *et al.*, 2019; Lilles *et al.*, 2015), healthy plants provide low and high reflectance values in red and near-infrared waves, and the NDVI is expressed in the range -1 to 1. Negative and positive values indicate non-vegetative and vegetative objects, where an NDVI value close to 1 indicates dense and healthy vegetation.

Nutrient Analysis Results

Soil analysis showed significant difference in the parameters of pH H₂O, C-Organic, P, and K. There was no difference in soil nitrogen content on land planted with oil palm aged 7, 12 and 16 years. The application of various fertilisers did not affect soil N content as shown in *Table 8*.

TABLE 6. DISTRIBUTION OF OIL PALM ROOT TO VARIOUS TYPES OF PLANTING AGES AND APPLICATIONS OF VARIOUS TYPES OF FERTILISER

Plant age (yr)	Types of fertiliser	Distribution of root					
		Primary		Secondary		Tertiary	
7	Control	0.41	ab	0.27	ab	0.16	ab
	K fertiliser	0.66	a	0.42	a	0.25	a
	Leaf litter	0.49	ab	0.34	ab	0.20	ab
	Empty bunches	0.51	ab	0.34	ab	0.20	ab
12	Control	0.31	b	0.22	ab	0.13	ab
	K fertiliser	0.49	ab	0.35	a	0.21	a
	Leaf litter	0.36	ab	0.28	ab	0.17	ab
	Empty bunches	0.38	ab	0.28	ab	0.17	ab
16	Control	0.13	c	0.10	c	0.05	c
	K fertiliser	0.19	c	0.14	bc	0.08	bc
	Leaf litter	0.16	c	0.11	c	0.07	c
	Empty bunches	0.16	c	0.11	c	0.06	c

TABLE 7. OIL PALM NDVI FOR VARIOUS TYPES OF PLANTING AGES AND APPLICATIONS OF VARIOUS TYPES OF FERTILISER

Plant age (yr)	Types of fertiliser	NDVI	
7	Control	0.736	A
	K fertiliser	0.754	A
	Leaf litter	0.737	A
	Empty bunches	0.756	A
12	Control	0.682	B
	K fertiliser	0.681	B
	Leaf litter	0.694	B
	Empty bunches	0.682	b
16	Control	0.682	b
	K fertiliser	0.681	b
	Leaf litter	0.694	b
	Empty bunches	0.682	b

Soil pH from *Table 8* showed significant difference for plants of different ages with the application of various types of fertiliser. Plants aged 7 years with the application of leaf litter and empty fruit bunches had the highest pH values, namely 5.65 and 6.00 (slightly acidic) and the lowest were found in plants aged 12 years with control treatments and application of K fertiliser (4.38 and 4.30 with very acidic criteria). The use of midrib and empty fruit bunches, both rich sources of organic matter, significantly influenced soil pH levels when subjected to decomposition. The breakdown of these organic materials generated compounds that altered the pH of the soil. Furthermore, the humification process stemming from coarse organic matter decomposing produces humic and fulvic acids, both of which increased soil pH. Butterley *et al.* (2013) affirmed that the use of dry matter in the form of residue, at a rate of

10 g/kg, was capable of augmenting soil pH values, leading to a temporary shift towards alkalinity. The condition was a consequence of the decomposition processes, driven by abiotic association reactions between H⁺ ions and the introduced organic matter. Additionally, the processes of ammonification and decarboxylation, occurring during decomposition, contributed to the generation of alkaline conditions in soil.

The application of leaf litter fertiliser and empty fruit bunches on plants aged 12 years increased the organic C content of the soil to around 9.72-10.3 (very high). The use of midrib litter and empty fruit bunches as a source of organic matter increased the availability of soil organic C when decomposed. The application of these two materials was also a form of using oil palm waste which was returned to the soil as a source of nutrients. Some results showed that the use of waste palm fronds and leaf residues restored soil nutrients and increased soil organic C (Ariyanti *et al.*, 2019; Singh *et al.*, 2013).

Table 8 shows that the soil P content of 16-year-old oil palms was much higher than 7 and 12-year-old plants for various types of fertiliser applications. However, the application of different types of fertiliser did not have a significant effect on the phosphorus content in oil palms aged 16 years for both controls, K fertiliser, midrib leaf litter, and empty fruit bunches with P values ranging from 144.19 to 152.45 (very high). This was due to the accumulation of P during fertilisation which was carried out until the age of 16 years. For soil K content there were also significant differences in the use of various types of fertiliser at various plant ages. The highest K value was found in the treatment of plant leaf litter aged 7 years after planting with the application of leaf litter to provide the needs in soil.

N content in soil at various ages of oil palm with the use of various types of fertiliser did not show significantly different results with N values ranging from 0.17 to 3.05 (low-very high). The nitrogen content that did not differ in soil was influenced by the availability of sufficient amounts in soil. This nutrient was the most important limiting factor for plant productivity, and its availability was very important (Niu *et al.*, 2016). Excessive use of fertiliser in agricultural management causes loss of N through leaching and denitrification processes (Liu *et al.*, 2010).

Root NPK levels based on analysis of variance reported significant differences in observed parameters both in N, P and K content. Table 9 showed that plant age and the application of various types of fertiliser had a significant effect on root NPK.

Table 9 shows the NPK levels in the roots of various ages of oil palms due to the application of fertiliser. Root N content in oil palms aged

12 and 16 years with the use of frond litter fertiliser showed the highest N content compared to others. The use of midrib litter was able to provide nitrogen in the roots of oil palms and was used for plant growth. Yama (2018) stated that midrib litter contained a C/N ratio of 16, Organic 34.18, N 2.09%, P 826 ppm, and K 1790 ppm. This litter was obtained from fallen fruit, fronds, and stems stored in the armpits of fronds and experienced weathering. Meanwhile, weathered organic matter enriched the planting medium, and the planting medium became fertile, loose, and grew freely. The results of other studies indicated that litter on oil palm stems was used to replace topsoil as a nursery medium for *Pueraria javanica* plants.

For the P content in the root, oil palms at the age of 12 years reported the highest content compared to the ages of 7 and 16 years using K fertiliser, leaf litter, and empty fruit bunches. Table 9 shows that the numerical content of leaf litter has a high value compared to K fertiliser and empty fruit bunches.

TABLE 8. OIL PALM NDVI FOR VARIOUS TYPES OF PLANTING AGES AND APPLICATIONS OF VARIOUS TYPES OF FERTILISER

Plant age (yr)	Types of fertiliser	Soil analysis									
		pH H ₂ O		C-org.		N		P		K	
7	Control	4.86	ab	2.12	b	0.30	a	24.75	b	0.65	b
	K fertiliser	4.68	ab	4.29	ab	0.31	a	46.25	ab	1.09	b
	Leaf litter	6.00	a	5.75	ab	0.31	a	55.81	ab	2.18	a
	Empty bunches	5.65	a	1.93	b	0.17	a	22.77	b	1.51	ab
12	Control	4.38	b	5.96	ab	0.30	a	97.36	ab	0.26	b
	K fertiliser	4.30	b	9.25	ab	0.41	a	103.59	ab	0.40	b
	Leaf litter	5.46	ab	10.3	a	0.34	a	120.71	ab	0.93	b
	Empty bunches	4.88	ab	9.72	a	0.30	a	101.83	ab	0.82	b
16	Control	4.95	ab	6.45	ab	2.76	a	145.49	a	0.37	b
	K fertiliser	4.80	ab	9.31	ab	0.50	a	152.45	a	0.34	b
	Leaf litter	5.15	ab	6.89	ab	3.05	a	147.06	a	0.40	b
	Empty bunches	4.70	ab	5.98	ab	0.63	a	144.19	a	0.45	b

TABLE 9. NPK CONTENT OF OIL PALM PLANT ROOT FOR VARIOUS TYPES OF PLANTING AGES AND THE APPLICATION OF VARIOUS TYPES OF FERTILISER

Plant age (yr)	Types of fertiliser	Root NPK levels					
		N		P		K	
7	Control	0.15	c	0.22	b	0.68	a
	K fertiliser	0.17	c	0.22	b	0.54	ab
	Leaf litter	0.19	bc	0.18	b	0.78	a
	Empty bunches	0.12	c	0.15	b	0.56	ab
12	Control	0.30	ab	0.22	b	0.31	b
	K fertiliser	0.21	ab	0.58	a	0.51	ab
	Leaf litter	0.38	a	0.68	a	0.70	a
	Empty bunches	0.20	ab	0.51	a	0.50	ab
16	Control	0.19	bc	0.27	b	0.58	ab
	K fertiliser	0.16	c	0.36	b	0.63	ab
	Leaf litter	0.35	a	0.31	b	0.66	ab
	Empty bunches	0.19	bc	0.23	b	0.24	b

This was because the leaf litter had a high P content and when applied as fertiliser more nutrient was provided to be absorbed. The application of empty fruit bunches of fertiliser to 12-year-old palms also provided high P. According to Hayat and Andayani (2014), the nutrient content of empty palm fruit bunches compost contained a total of N (1.91%), K (1.51%) and Ca (0.83 %), P (0.54%), Mg (0.09%), C-organic (51.23%), C/N ratio 26.82% and pH 7.

The application of K fertiliser indirectly increased the P content in the root of oil palms. The use provides potassium to plants and indirectly affects the absorption of P in the root. Potassium fertiliser given correctly can increase the efficiency of P use by plants, and assist in the process of transporting the element in plants to increase the efficiency of absorption. The element improves the quality of plant roots and explores more areas of soil, including areas rich in P.

CONCLUSION

In conclusion, the growth of plants aged 7 years was the best response compared to plants aged 12 and 16 years in terms of stem circumference, stem diameter, and NDVI values for all types of fertiliser applied. The primary root distribution for plants aged 7 years with the use of K fertiliser showed the best results for the distribution of secondary and tertiary roots and was not significantly different from those aged 12 years. Soil analysis showed that there were differences in the use of various types of fertiliser on soil nutrient content, specifically on pH, C-organic, P, and K values. The use of midrib and empty fruit bunches affected plants aged 7 years, while those aged 12 years had C content. The high organic matter with the use of leaf litter and empty fruit bunches for oil palms aged 16 years had the best effect on P values compared to those aged 7 and 12 years.

ACKNOWLEDGEMENT

The authors are grateful to the Indonesian Endowment Fund for Education (LPDP) 2019 (0007721/TRP/D/BUDI-2019), Ministry of Finance, Indonesia for their financial support.

REFERENCES

Amri, A I; Armaini, A and Amindo Purba, M R (2018). Application of empty palm fruit bunch compost and dolomite on inceptisol subsoil medium to oil palm (*Elaeis guineensis* Jacq.) seedlings in main nurseries. *J. Agroteknol.*, 8(2): 1. DOI: 10.24014/ja.v8i2.3349.

Ariyanti, M; Maxiselly, Y; Rosniawaty, S and Indrawan, R A (2019). Pertumbuhan kelapa sawit belum menghasilkan dengan pemberian pupuk organik asal pelepah kelapa sawit dan asam humat. *Jurnal Penelitian Kelapa Sawit*, 27(2): 71-82.

Baryanto; Nelvia and Wardati (2015). Pengaruh pemberian kompos tandan kosong kelapa sawit (TKSS) pada pertumbuhan bibit kelapa sawit (*Elaeis guineensis* Jacq.) di main-nursery pada medium subsoil ultisol [The effect of composting empty palm oil bunches (EFB) on the growth of oil palm seedlings (*Elaeis guineensis* Jacq.) in main-nursery on subsoil ultisol medium. *JOM.*, 2(1).

Budiarto, R; Poerwanto, R; Santosa, E and Efendi, D (2019). A review of root pruning to regulate citrus growth. *J. Trop. Crop Sci.*, 6(01): 1-7. DOI: 10.29244/jtcs.6.01.1-7.

Butterly, C R; Baldock, J A and Tang, C (2013). The contribution of crop residues to changes in soil pH under field conditions. *Plant Soil*, 366: 185-198. DOI: 10.1007/s11104-012-1422-1.

Cahyono, B E; Febriawan, E B and Nugroho, A T (2019). Land cover analysis using landsat imagery unguided classification method in Sawahlunto, West Sumatra. *Jurnal Teknotan*, 13(1): 8-14. DOI: 10.24198/jt.vol13n1.2.

Cha-um, S; Samphumphuang, T and Kirdmanee, C (2013). Glycinebetaine alleviates water deficit stress in indica rice using proline accumulation, photosynthetic efficiencies, growth performances and yield attributes. *Aust. J. Crop Sci.*, 7(2): 213-218.

Endriani; Zurhalena and Refliaty (2003). Improvement of ultisol soil physical properties and crop yields through the application of bokashi fertilizer. *Prosiding Buku I. Kongres Nasional VIII Himpunan Ilmu Tanah Indonesia*. Padang, Indonesia. 21-23 Juli 2003.

Fangeria, N K; Filho, M P B and Da Costa, J H C (2009). *Potassium in the Use of Nutrients in Crop Plants*. CRC Press Taylor & Francis Group, Boca Raton, London, New York. p. 131-163.

Fini, A; Frangi, P; Faoro, M; Piatti, R; Amoroso, G and Ferrini, F (2015). Effects of different pruning methods on an urban tree species: A four-year-experiment scaling down from the whole tree to the chloroplasts. *Urban For. Urban Green*, 14(3): 664-674.

Foong, A; Sum, W and Shukor, S A A (2019). Oil palm plantation monitoring from satellite image

- recent citations a new machine learning approach in detecting the oil palm plantations using remote sensing data kaibin oil palm plantation monitoring from satellite image. *IOP Conf. Ser. Mat. Sci. Engin.*, 705(0): 2.
- Habi, M L (2015). Effect of application of ela sago granule compost enriched with Ponska fertilizer on soil physical properties and yield of sweet corn in Inceptisols. *Biopendix*, 1(2): 121-134.
- Hayat, E S and Andayani, S (2014). Management of empty oil palm fruit bunches and the application of chromolaena odorata biomass on growth and yield of rice plants and sulfaquent soil properties. *Jurnal Teknologi Pengelolaan Limbah*, 17(22): 44-51.
- Kassem, H A and Marzouk, H A (2010). Improving yield and fruit quality of date palm by organic fertilizer sources. *J. Appl. Horticult.*, 12(2): 145-150.
- Lillesand, T M; Kiefer, R W and Chipman (2015). *Remote Sensing and Image Interpretation*. Wiley. 736 pp.
- Liu, J; You, L and Amini M *et al.* (2010). A high-resolution assessment on global nitrogen flows in cropland. *Proc. Natl. Acad. Sci. USA*, 107: 8035-8040. DOI: 10.1073/pnas.0913658107.
- Luok, M and Raharjo, K T P (2017). The effect of root cutting and acclimatization time on the growth of candlenut plant seedlings (*Aleurites moluccana* Willd) from Stum. *Savana Cendana*, 2(02): 19-22. DOI: 10.32938/sc.v2i02.87.
- Niu, S; Classen, T; Dukes, J S; Kardol, P; Liu, L; Luo, Y; Rustad, L; Sun, J; Tang, J; Templer, P H *et al.*, (2016). Global patterns and substrate-based mechanisms of the terrestrial nitrogen cycle. *Ecol. Lett.*, 19: 697-709. DOI: 10.1111/ele.12591.
- Nurcahyo, N E and Ariani, E (2017). The effect of gandasil fertilizer and a variety of waste oil palm plantation on the growth of seedlings of oil (*Elaeis guineensis* Jacq.) seedlings in the main nursery. *JOM Faperta*, 14(1): 1-13.
- Miller, F D and Neely, D (1993). The effect of trenching on growth and plant health of selected species of shade trees. *J. Arboric.*, 19: 226-229.
- Rao, I M; Borrero, V; Ricaurte, J and Garcia, R (1999). Adaptive attributes of tropical forage species to acid soils: Differences in phosphorus acquisition from inorganic and organic phosphorus sources. *J. Plant Nutr.*, 22: 1175-1196. DOI: 10.1080/01904169909365703.
- Rozy, F; Rosmawaty, T and Fatrrahman (2013). Pemberian pupuk N P K Mutiara 16:16:16 dan kompos tandan kosong kelapa sawit pada tanaman terung (*Solanum melongena* L). *Jurnal RAT.*, 1(2): 228-239.
- Sarwono, E (2008). Utilization of empty buds as a substitute for fertilizer for palm oil plants. *APLIKA.*, 8(1).
- Singh, P; Sulaiman, O; Hashim, R; Peng, L C and Singh, R P (2013). Evaluating biopulping as an alternative application on oil palm trunk using the whiterot fungus *Trametes versicolor*. *Int. Biodeter. Biodegrad.*, 82: 96-103. DOI: 10.1016/j.ibiod.2012.12.016.
- Syakir, M and Gusmaini (2012). The effect of using potassium fertilizer sources on the production and quality of patchouli oil. *Jurnal Littri*, 18(2): 60-65.
- Watson, G W (1998). Tree growth after trenching and compensatory crown pruning. *J. Arboricult.*, 24(1): 47-53.
- Widodo, K H; Kusuma, Z; Tanah, J; Pertanian, F; Brawijaya, U and Korespondensi, P (2018). Effect of compost on soil physical properties and growth of corn plants in inceptisols. *J. Tanah dan Sumberd. Lahan*, 5(2): 959-967.
- Greenland, D (2003). *Sustainable Soils: The Place of Organic Matter in Sustaining Soils and Their Productivity* (Wolf, B and Snyder, G H eds.). New York: Haworth Press. 352 pp.
- Yahya, Z; Husin, A; Talib, J; Othman, J; Ahmed, O H and Jalloh, M B (2010). Oil palm (*Elaeis guineensis*) roots response to mechanization in Bernam series soil. *Am. J. Appl. Sci.*, 7(3): 343- 348. DOI: 10.3844/ajassp.2010.343.348.
- Yama, D I (2018). Analisis pertumbuhan pembibitan *Pueraria javanica* pada komposisi media seresah dalam ketiak pelepah pada batang kelapa Sawit. *Jurnal Citra Widya Edukasi*, 10(3): 199-206.
- Yuniati, S and Widyastuti, R (2014). *Composting palm oil midrib-leaf with different biodecomposters and used as ameliorant*. Master thesis. Bogor: Institut Pertanian Bogor.
- Yuniasih, B; Adj, A R P and Budi, B (2022). Evaluation of pre-replanting oil palm plant health using the NDVI index from Landsat 8 satellite imagery. *J. Agric. Engin.*, 11(2): 304-313. DOI: 10.23960/jtep-l.v11i2.304-313.