

APPLICATION OF OLD OIL PALM TRUNKS AFFECTS THE GROWTH PERFORMANCE OF OIL PALM SEEDLINGS

AYAKA UKE^{1*}; ALISA HANIS²; KUMAR SUDESH² and AKIHIKO KOSUGI^{1,3*}

ABSTRACT

Oil palm plantations generate substantial biomass waste, primarily old oil palm trunks (OPT), during replanting. As part of a sustainable plantation management program, old OPT are returned to the plantation and released nutrients into the soil for new oil palm seedlings. However, whether this method improves soil nutrient levels is unclear. In this study, we investigated the effects of OPT on oil palm seedling growth and soil microbial communities. The plant height, chlorophyll content, leaf area, and biomass weight were low in seedlings grown in soil containing OPT (44.3 cm, 44,278.9 cm², 19.6 g [dry shoot], and 14.9 g [dry root]). Similar results were obtained for seedlings grown in soil containing cellulose or soil containing OPT and fertiliser. Leaf nitrogen, phosphorus, and potassium contents were similar in seedlings grown in soil amended with OPT and control seedlings. However, the calcium content was significantly lower in seedlings grown in soil containing OPT (0.424 ± 0.004%) than in control seedlings (0.496 ± 0.006%). Metagenomic analysis of soils showed that three lignocellulose-degrading fungal genera (*Chaetomium*, *Mortierella*, and *Staphylotrichum*) were abundant in soil containing OPT. Thus, the return of OPT promotes the growth of lignocellulose-degrading microorganisms and decreases fertiliser nutrient availability.

Keywords: *Chaetomium*, lignocellulose-degrading microorganisms, oil palm trunk, plant growth performance, soil metagenomics.

Received: 18 May 2024; **Accepted:** 26 November 2024; **Published online:** 20 February 2025.

INTRODUCTION

Palm oil is regarded as one of Malaysia's primary sector industries and is a major agricultural product exported globally. Oil palm productivity decreases every 20 to 25 years, and new oil palm seedlings are replanted (Corley & Tinker, 2015). The old oil palm trees are felled to clear the land, and are left in the plantation to decompose. Some

plantations use the chip and windrow method (e.g., Guan Soon Plantation, Alor Pongsu, Perak, Malaysia), whereas some use the pulverisation technique, such as Kuala Lumpur Kepong Berhad (2018), whereby entire old oil palm trunks (OPT) are pulverised into smaller pieces that are then spread on the cleared land (Pulingam et al., 2022). OPT left using the windrow method takes at least two years to decompose completely, whereas those that are pulverised and spread onto soil degrade within approximately one year, which helps to shorten the fallow period. The decomposing old OPT is considered to serve as mulch (Sung, 2016) while simultaneously replenishing soil nutrients as they break down (Pulingam et al., 2022).

However, during the decomposition period, various environmental issues may arise. For example, OPT eventually become a breeding site for pests, such as *Oryctes rhinoceros* (Manjeri et al., 2014), and the fungus *Ganoderma boninense*, which causes basal stem rot disease (Gorea et al., 2019).

¹ Biological Resources and Post-Harvest Division, Japan International Research Center for Agricultural Sciences (JIRCAS), 1-1 Ohwashi, Tsukuba, Ibaraki 305-8686, Japan.

² Ecobiomaterial Research Laboratory, School of Biological Sciences, Universiti Sains Malaysia, 11800 Pulau Pinang, Malaysia.

³ University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8577, Japan.

* Corresponding author e-mail: ukea0222@jircas.go.jp, kosugia0164@jircas.go.jp

Additionally, the high carbon-to-nitrogen (C:N) ratio (Lin et al., 2019) lengthens the decomposition period (Akratos et al., 2017) and deprives the soil of nitrogen, which is needed for plant growth (Lin et al., 2019). Usually, a C:N ratio between 25 and 35 is optimal for agricultural applications; however, OPT has a high C:N ratio of 155 (Loh et al., 2013) and it takes a long time for this ratio to decrease. Therefore, the substantial OPT biomass waste generated during replanting and left in oil palm plantations to decompose may negatively affect plant growth. The application of OPT results in decreased growth performance compared with empty fruit bunches (Mohammed et al., 2014). It also alters the soil microflora and decreases the availability of nutrients, such as nitrogen, calcium and magnesium, for plant uptake (Uke et al., 2021). Nitrogen promotes leaf growth. The leaf is one of the most important plant organs because it is a nitrogen storage site that synthesises amino acids. Calcium is an important component of cell walls and membranes, thereby contributing to the structural integrity of cells. This helps to maintain the physical barriers that protect plants from pathogens, leading to enhanced immunity. Magnesium is a component of chlorophyll, which is essential for photosynthesis. A magnesium deficiency may cause chlorosis.

The soil type and quality of organic matter affect the diversity and abundance of soil microflora (Li et al., 2018; Savy et al., 2020). The soil and rhizosphere microorganisms are bio-indicators of soil quality because they are sensitive to the smallest changes in abiotic conditions such as environmental stress and soil perturbation. In addition, land fertility and environmental biodiversity are affected by the cultivation of oil palm (Ashton-Butt et al., 2018; McGuire et al., 2014).

To understand how OPT breakdown affects soil health and microbial diversity within a plantation, it is important to determine the effect of applying OPT fibre directly to the soil, mimicking the pulverisation technique, on oil palm seedling growth and the soil microbial community. In this study, we explored the effects of OPT fibre and the contrasting effects of fertiliser applications on the growth of oil palm seedlings. We experimented with various soil amendments: Unamended soil (T1), soil + OPT fibre (T2), soil + cellulose (T3), soil + fertiliser (T4), soil + OPT fibre + fertiliser (T5), and soil + cellulose + fertiliser (T6). Soil metagenomics analyses were conducted to explore the diversity and abundance of soil microflora. Thus, this study provides valuable insights into the sustainability and productivity of oil palm plantations by determining the effects of OPT biomass waste on the growth of oil palm seedlings and the soil microbial community.

MATERIALS AND METHODS

Polybag Assays

To establish the different treatments, dried OPT fibre, cellulose powder as cellulosic material, and fertiliser was added to the soil that was collected from an Malaysian Palm Oil Board (MPOB)-certified oil palm nursery in Alor Pongsu, Perak, Malaysia. Fresh OPT fibre (>20 years old) was air-dried under the sun until the moisture content reached <10% (w/w) to prevent any microbial growth (Lai et al., 2014). Fifty-five 4-month-old oil palm seedlings (*Elaeis guineensis*) were used for the polybag assay, which was conducted in an open space at an oil palm seedling distributor in Alor Pongsu (5°04'37.8" N, 100°35'52.4" E). Twin Arrow Fertiliser (TAF) (15:15:6:4/N:P:K:Mg+TE), a commercial fertiliser from TAF Sdn. Bhd., Selangor, Malaysia and cellulose fibre powder (Arbocel®, Germany) were used. The polybag assay was further modified according to our previous study (Uke et al., 2021). The four-month-old seedlings were transplanted into polybags (33 × 35 cm) containing 10 kg soil. The six treatments were prepared as shown in Table 1. In total, 55 polybags of oil palm seedlings were prepared, with 10 replicates for Treatment 1, 2, 4 and 5; seven replicates for Treatment 3; and eight replicates for Treatment 6. All polybags containing seedlings were arranged in rows with spacing of 0.75 × 0.75 × 0.75 m following the "Code of Good Nursery Practice for Oil Palm Nurseries" published by the MPOB (2016). A standardised average of at least seven replicates of oil palm seedlings per treatment was used to validate the results. The treatments were applied using a factorial and randomised design. The amount of OPT fibre added to each polybag (200 g) was determined using Equation (1). The OPT fibre was added to the soil and mixed thoroughly. This was based on an initial dry weight of OPT of approximately 1,315 kg. However, the amount was reduced by half to ensure that the OPT fibre could fit in each polybag. Seedlings were grown under a natural photoperiod and were watered daily. Fertiliser was added to certain treatments of oil palm seedlings (Treatment 4, 5 and 6) at the dose recommended by the manufacturer at the start of the experiment.

$$\text{Dry weight of OPT} \times \text{Number of oil palm trees per hectare} \times \text{Area of polybag} \quad (1)$$

Plant Measurements

The plant height, relative leaf area, relative chlorophyll content (as determined using a Soil

TABLE 1. TREATMENTS APPLIED TO OIL PALM SEEDLINGS OVER A FOUR-MONTH GROWTH PERIOD

Treatment label	Treatments	Amounts of treatments applied (g/polybag)
T1	Unamended soil (control)	No amendments
T2	Oil palm trunk (OPT) fibre	200 g
T3	Cellulose powder	200 g
T4	Fertiliser	7 g
T5	OPT fibre + Fertiliser	200 g of OPT fibre + 7 g fertiliser
T6	Cellulose + Fertiliser	200 g of cellulose + 7 g fertiliser

Plant Analysis Development [SPAD] meter), soil pH, and biomass weight were measured to determine the growth performance of oil palm seedlings in the various treatments (Uke et al., 2021). The plant height was measured from the ground level up to the top shoot (Adip et al., 2022). The relative leaf area was determined by multiplying the length and width of the fully extended leaves (length × width). The relative chlorophyll content was measured using a Minolta SPAD-520 Plus meter (Konica Minolta Inc., Japan). A soil pH and moisture tester (Model DM-15, Takemura Electric Works Ltd., Japan) was used to determine the soil pH. Measurements were taken at the start of the experiment and every month for four consecutive months. The biomass dry weight was determined at the end of the 4-month growth period by oven-drying the shoots and roots to a constant weight at 70°C.

Leaf Nutrient Analysis

The leaf samples were analysed by Applied Agricultural Resources Sdn. Bhd., Malaysia. The nitrogen, phosphorus, potassium, calcium, and magnesium contents of oil palm seedling leaves were determined. Kjeldahl's method was used to analyse the total nitrogen content and the acid digestion method was used to analyse the phosphorus content. The potassium content was determined using the ammonium acetate leaching method followed by atomic absorption spectrophotometry using a Perkin Elmer Analyst 100 Atomic Absorption Spectrometer (USA).

High-throughput Sequencing

DNA extraction. Soil from each treatment was used for DNA extraction. Before soil samples were collected, 1 to 2 cm of the top soil layer was removed. For each treatment, soil was collected through mixing. The samples were immediately stored at -20°C until DNA extraction was conducted. Total genomic DNA was extracted from 250 mg soil using the DNeasy PowerSoil Pro Kit (Qiagen, , Germany) according to the manufacturer's instructions. The DNA concentration was determined

using a NanoDrop One Microvolume UV-VIS Spectrophotometer (Thermo Fisher Scientific, USA). A polymerase chain reaction (PCR) targeting the internal transcribed spacer 1 (ITS) region of the fungal rRNA gene was conducted using the primers ITS1F_KYO1 (CTHGGTCATTTAGAGGAASTAA) and ITS2_KYO2 (TTYRCTRCTTCATC) as well as Tks Gflex DNA Polymerase Low DNA (TaKaRa, Japan). The DNA amplification conditions were as follows: Initial denaturation at 94°C for 1 min; 35 cycles of 98°C for 10 s, 55°C for 15 s, and 68°C for 1 min; and final extension at 68°C for 5 min. The reaction products were separated on a 2% (w/v) agarose gel, stained with ethidium bromide, and purified (Ungkulpasvich et al., 2021). After diluting to 25 pM, all sample libraries were combined for an automated emulsion PCR and bead enrichment and then loaded onto 510TM, 520TM, and 530TM chips for sequencing using an Ion ChefTM instrument (Thermo Fisher Scientific). The libraries were sequenced using the Ion GeneStudioTM S5 System with 850 flow cycles.

Operational taxonomic unit clustering and community analyses. The raw sequences generated by the Ion ChefTM instrument were subjected to further filtering and trimming to obtain sequences >150 bp and <300 bp using CLC Genomics Workbench v.23.0.4 and CLC Microbial Genomics Module v.20.1 (Qiagen, USA). The UNITE database was used to identify the operational taxonomic units (OTUs) with 97% nucleotide identity (Uke et al., 2021). The software automatically identified and discarded chimeric sequences. The filtered read sets were then grouped into OTUs with 0.02 distance unit cut-offs according to the BLASTN algorithm in the National Center for Biotechnology Information GenBank database. The abundance and diversity of the fungi present in the soil were analysed.

Statistical analysis. Statistical analyses were conducted using SPSS (version 24.0) (IBM Inc., USA). Data were subjected to a one-way ANOVA, with means separated using the least significant difference test. Differences were considered significant at $P \leq 0.05$

RESULTS AND DISCUSSION

Effects of Adding Oil Palm Trunk (OPT) Fibre to Soil on Plant Growth Performance

The effects of adding OPT fibre and other substances to soil on plant growth performance (i.e., plant height, relative leaf area, chlorophyll content and biomass) were determined after four months of growth (*Figure 1*). At the end of the 4th month after application (MAA), the height of oil palm seedlings was significantly lower in the OPT fibre (T2) and cellulose treatments (T3) than in the unamended (T1) and fertiliser treatments (T4). Moreover, the growth indexes of seedlings in T2 and T3 were significantly different from those of seedlings in the T1, T4, OPT + fertiliser (T5) and cellulose + fertiliser (T6) treatments. This implied that fertiliser nutrient availability decreased in the presence of OPT fibre and cellulose. Interestingly, the comparison with the T1 revealed plant growth and development were significantly inhibited in the T2 and T3 treatments. Similarly, the relative leaf area and the chlorophyll content of the oil palm seedlings were significantly lower in T2 and T3 treatments than in the other treatments and T1 (*Figure 1b* and *1c*).

The small relative leaf area, homogenous yellowing of leaves, and stunted growth observed in T2 and T3 are symptoms of chlorosis. The lower height of the oil palm seedlings in these treatments, compared with that in T1, was associated with decreases in the chlorophyll content and leaf area. The chlorophyll content (Wen et al., 2019), total nitrogen content, soluble protein content in leaves, and the net photosynthetic rate are all affected by nitrogen availability (Qu et al., 2022). Plants under nitrogen deficiency stress have a reduced photosynthetic capacity (Qu et al., 2022), which eventually leads to poor growth and root development (Kang et al., 2023). Our results also show that the seedlings in T2 and T3 had less fibrous roots than T1 (*Figure 2*). It has been reported that the morphological development of oil palm roots (length, surface area, and volume) is restricted under nitrogen-deficient conditions (De la Peña et al., 2024).

The reduced plant biomass (both shoot and root) in T2 and T3 may be because of microbial nitrogen immobilisation (Uke et al., 2021). The carbon content affects the amount of nitrogen immobilised by the soil microbial community. Microbial biomass and respiration are relatively low under high C:N conditions because of the limited availability of nitrogen for microorganisms, leading to delayed degradation of plant fibre. Crop residues with high C:N ratios include OPT fibre (155) (Loh et al., 2013) and sugarcane bagasse (213) (Bhat et al., 2015). In another study, the application of sugarcane straw affected the growth and development of sugarcane plants, although the yield and quality of

sugarcane juice were unaffected (Souza et al., 2020). Our results show that OPT fibre left in the soil eventually decreased the efficiency of soil microbe-mediated degradation of cellulosic biomass waste because of its high C:N ratio. Additionally, it takes approximately two years for OPT to decompose and the high C:N ratio of OPT is expected to decrease during the degradation period. This finding highlights the effects of OPT degradation in soil. Although this study was conducted using polybags, similar effects may be observed in oil palm plantations.


Effects of Various Treatments on Nutrient Contents in Oil Palm Seedling Leaves

The nitrogen content in oil palm seedling leaves was higher in T2 (2.19 ± 0.01) and T5 (3.41 ± 0.008) than in T1 (1.94 ± 0.01). The calcium content in oil palm seedling leaves was higher in T1 (0.50 ± 0.006) than T2 and T5 (0.42 ± 0.004 and 0.32 ± 0.002 , respectively) and the magnesium content was also higher in T1 (0.46 ± 0.004) than in T2 and T5 (0.43 ± 0.005 and 0.38 ± 0.002 , respectively) (*Table 2*). These results show that the addition of fertiliser to the soil amended with OPT fibre did not increase the availability of nutrients for new oil palm seedlings.

According to the results of the foliar analysis of oil palm seedlings, we speculated that the OPT fibre may have negatively affected leaf growth because of the low contents of certain nutrients, including nitrogen, magnesium, and calcium. This is supported by the overall plant growth parameters (i.e., height, chlorophyll content, relative leaf area, and shoot and root biomass) that were adversely affected. Similarly, a previous study showed the addition of wheat residue to soil (applied to the soil surface or mixed with the soil) delayed winter wheat seed emergence; the resulting seedlings were shorter than normal and exhibited an abnormal geotropic response, which may reflect the phytotoxic effect of the residue (Wuest et al., 2000). The application of winter wheat residue is similar to the application of OPT in oil palm plantations using the chip and windrow method and the pulverisation technique. Thus, to replenish nitrogen losses in soil, larger amounts of nitrogen fertiliser were required for the new plants to grow and achieve a satisfactory yield.

Effects of Various Treatments on the Soil Microbial Community

Plant growth is affected by soil fertility, which in turn is affected by the soil microbial population. The structure of soil microbial communities determines the quality of soil in oil palm plantations because microbes function as carbon recycling agents






Figure 1. Growth performance of oil palm seedlings in soil containing OPT fibre, cellulose, fertiliser, OPT fibre + fertiliser, and cellulose + fertiliser. (a) Height, (b) leaf area, (c) chlorophyll content, and (d) dry weight of the oil palm seedling shoot and root. Significant differences were observed among the treatments at 4 MAA (ANOVA; $F = 22.8$; $P < 0.001$ for height, ANOVA; $F = 16.9$; $P < 0.001$ for relative leaf area, ANOVA; $F = 41.1$; $P < 0.001$ for chlorophyll content, ANOVA; $F = 16.8$; $P < 0.001$ for shoot dry weight, ANOVA; $F = 8.591$; $P < 0.001$ for root dry weight). $N = 10$ for control, OPT fibre, fertiliser, and OPT fibre + fertiliser, $N = 7$ for cellulose only, and $N = 8$ for cellulose + fertiliser. Significant differences among treatments at specific time points are indicated with asterisks (** $P < 0.001$).




Figure 2. (a) Growth of oil palm seedlings in various treatments and (b) roots and above-ground parts of oil palm seedlings. Treatment 1: Unamended soil (control), Treatment 2: OPT fibre (200 g), Treatment 3: Cellulose (200 g), Treatment 4: Fertiliser (7 g), Treatment 5: OPT fibre (200 g) + fertiliser (7 g), and Treatment 6: Cellulose (200 g) + fertiliser (7 g).

TABLE 2. NUTRIENT CONTENTS IN OIL PALM LEAVES

Elements	Unamended soil (T1) (%)	OPT fibre (T2) (%)	OPT fibre + fertiliser (T5) (%)
Nitrogen (N)	1.942 ± 0.010 ^e	2.188 ± 0.010 ^d	3.410 ± 0.008 ^c
Phosphorus (P)	0.205 ± 0.001 ^c	0.227 ± 0.001 ^b	0.241 ± 0.0003 ^a
Potassium (K)	1.918 ± 0.008 ^e	2.398 ± 0.030 ^{ab}	2.294 ± 0.010 ^c
Calcium (Ca)	0.496 ± 0.006 ^a	0.424 ± 0.004 ^b	0.316 ± 0.002 ^d
Magnesium (Mg)	0.460 ± 0.004 ^d	0.432 ± 0.005 ^d	0.382 ± 0.002 ^b

Note: OPT - oil palm trunk.

and as a source of nutrients (Situmorang et al., 2016). Microbes that function as carbon recycling agents include bacteria and fungi. Bacteria are active decomposers in soil and are responsible for decomposing dead plants and animals. Fungi are the predominant microbes in the soil microbial community (in terms of biomass) and they metabolise carbon-rich substrates (Wu et al., 2024).

Therefore, we analysed the eukaryotic microbial community structure in all the soil treatments (T1–T6). We focused on eukaryotic microbes because analyses of the ITS1 sequence data suggested that fungal OTUs showed the widest variation among treatments. The fungal genera *Chaetomium*, *Mortierella*, and *Staphylotrichum* were most abundant in T2 and T5 (Table 3).

TABLE 3. RELATIVE ABUNDANCE OF MICROBIAL SEQUENCES IN DIFFERENT SOIL AMENDMENT TREATMENTS

Abundance	T1.1	T1.2	T2	T3	T4	T5	T6
<i>Chaetomium</i> spp.	484	1,222	6,050	111	109	23,095	2,465
<i>Mortierella</i> spp.	4,756	4,224	1,445	296	2,549	1,076	91
<i>Staphylotrichum</i> spp.	0	2	4,497	0	0	601	0

Note: T1.1 - unamended soil at 0 MAA; T1.2 - unamended soil at 4 MAA; T2 - oil palm trunk (OPT) only; T3 - cellulose only; T4 - fertiliser only; T5 - OPT + fertiliser; T6 - cellulose + fertiliser.

This observation implies the application of OPT fibre results in a unique soil microflora community composition.

Chaetomium spp. in the Chaetomiaceae family has been found in soils amended with compost (Zhang et al., 2017). Members of this genus contain genes encoding enzymes that degrade lignocellulose biomass (Banerjee et al., 2016). *Chaetomium* spp. thrive and reproduce in the presence of lignin (Dicko et al., 2020), which is one of the major components of OPT. Additionally, *Mortierella* spp. accumulate lipids that are produced from sugars that are used as a carbon source (Ruan et al., 2012). OPT (70% moisture) have a high sugar content, especially during its decomposition period (Hanis et al., 2024; Yamada et al., 2010). Therefore, it can be inferred that the high relative abundance of these genera in the soil microflora was because the decaying OPT served as their main carbon source. Despite the abundance of *Staphylotrichum* spp. in the microflora in soil amended with OPT, to date, few studies have explored their role as lignocellulose-degrading fungi. However, one of the species in this genus, *Staphylotrichum longicolleum*, can degrade chitin in wood, sugarcane bagasse, and maize leaves (Ali et al., 2021).

During the decomposition of OPT, the starch in parenchyma cells is degraded and further fermented into sugars by wild endophytic fungi within the trunk (Abdul-Hamid et al., 2015). The high starch and sugar contents (i.e., glucose, fructose, and sucrose) of OPT (Hanis et al., 2024; Yamada et al., 2010) serve as the energy source for the growth and metabolism of microorganisms. Additionally, OPT has a high C:N ratio (Loh et al., 2013) and contains a considerable amount of xylan and cellulose (Uke et al., 2021), which influence and promote the accumulation of lignocellulose-degrading fungi. Thus, the composition of the soil microbial community may be affected by OPT degradation. All of the events that occur in the presence of OPT (De Lima Brossi et al., 2016) increase the abundance and diversity of certain soil microflora; this may result in the decreased availability of nutrients for plants, which was previously reported for corn plants grown in soil amended with OPT residue (Uke et al., 2021). According to soil metagenomics data, *Chaetomium* spp., *Mortierella* spp., and

Staphylotrichum spp. substantially affect nutrient dynamics. These microorganisms were highly effective at capturing and using available soil nutrients for their growth processes, which further decreased the availability of these nutrients for plants. The ability of these microorganisms to immobilise nutrients suggests that their presence and abundance may significantly influence soil nutrient profiles and, subsequently, plant productivity. These findings were results of previous studies, but they provide specific insights into the microbial species that predominantly affect nutrient availability in the current study.

CONCLUSION

Based on our results, we conclude that leaving the old OPT fibre residue in the plantation soil negatively affects the growth of oil palm seedlings because of soil nutrient deficiencies. Our results indicate that new oil palm seedlings were directly affected and grew relatively poorly (i.e., decreased height, leaf area, chlorophyll content and biomass) because of competition for nutrients with abundant lignocellulose-degrading fungi. In addition, fertiliser nutrient availability decreased in the presence of OPT. Thus, it is highly recommended that OPT be removed from oil palm plantations, even in small quantities, to decrease the amount of fertiliser required and to reduce fertiliser waste. The current practice of leaving OPT in the plantation requires a supplemental fertiliser application to compensate for the deficiency in available nutrients during the decomposition of OPT. To ensure a sustainable oil palm industry, the large amount of biomass in OPT should be used to its utmost potential for the production of value-added products (i.e., bioethanol, bio-pellets, and bioplastics).

ACKNOWLEDGEMENT

The authors thank the Ministry of Higher Education Malaysia (203/PBIOLOGI/67811001) for financial support and Edanz (<https://jp.edanz.com/ac>) for editing a draft of this manuscript.

This study was conducted as part of a development project funded by the Science and Technology Research Partnership for Sustainable Development (SATREPS) (Grant No. JPMJSA1801) of the Japan Science and Technology Agency (JST)-Japan International Cooperation Agency (JICA). The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this article.

REFERENCES

- Abdul-Hamid, Z., Arai, T., Mhd Ramle, S., Kosugi, A., Sulaiman, O., Hashim, R., Nirasawa, S., Ryohei, T., Eswara, L., Sudesh, K., Murata, Y., Saito, M., & Mori, Y. (2015). Analysis of free sugar and starch in oil palm trunks (*Elaeis guineensis* Jacq.) from various cultivars as a feedstock for bioethanol production. *International Journal of Green Energy*, 12(7), 635–641. <https://doi.org/10.1080/15435075.2014.910786>
- Adip, M., Meringgai, J., Aziz, N., Lah, M., & Yusoff, N. (2022). Growth performance of oil palm seedlings on different types of organic fertiliser. *Journal of Agricultural Biotechnology*, 13(2), 65–75. <https://doi.org/10.37231/jab.2022.13.2.325>
- Akratos, C. S., Tekerlekopoulou, A. G., Vasiliadou, I. A., & Vayenas, D. V. (2017). Composting of olive mill waste for the production of soil amendments. In C. M. Galanakis (Ed.), *Olive mill waste* (pp. 161–182). Academic Press.
- Ali, A., Ellinger, B., Brandt, S. C., Betzel, C., Rühl, M., Wrenger, C., Schlüter, H., Schäfer, W., Brognaro, H., & Gand, M. (2021). Genome and secretome analysis of *Staphylotrichum longicolleum* DSM105789 cultured on agro-residual and chitinous biomass. *Microorganisms*, 9(8), 1581. <https://doi.org/10.3390/microorganisms9081581>
- Ashton-Butt, A., Aryawan, A., Hood, A., Naim, M., Purnomo, D., Suhardi, S., Wahyuningsih, R., Willcock, S., Poppy, G., Caliman, J. P., Turner, E., Foster, W., Peh, K., & Snaddon, J. (2018). Understorey vegetation in oil palm plantations benefits soil biodiversity and decomposition rates. *Frontiers in Forests and Global Change*, 1, 10. <https://doi.org/10.3389/ffgc.2018.00010>
- Banerjee, S., Kirkby, C. A., Schmutter, D., Bissett, A., Kirkegaard, J. A., & Richardson, A. E. (2016). Network analysis reveals functional redundancy and keystone taxa amongst bacterial and fungal communities during organic matter decomposition in an arable soil. *Soil Biology and Biochemistry*, 97, 188–198. <https://doi.org/10.1016/j.soilbio.2016.03.017>
- Bhat, S. A., Singh, J., & Vig, A. P. (2015). Potential utilization of bagasse as feed material for earthworm *Eisenia fetida* and production of vermicompost. *SpringerPlus*, 4(1), 11. <https://doi.org/10.1186/s40064-014-0780-y>
- Corley, R. H. V., & Tinker, P. B. (2015). *The oil palm* (5th ed.). Wiley-Blackwell.
- De la Peña, M., Ruiz-Romero, R., Castro-Arza, L. I., & Romero, H. M. (2024). Changes in the root architecture of oil palm seedlings in response to nitrogen starvation. *Journal of Agronomy*, 14(3), 409–420. <https://doi.org/10.3390/agronomy14030409>
- De Lima Brossi, M. J., Jiménez, D. J., Cortes-Tolalpa, L., & van Elsas, J. D. (2016). Soil-derived microbial consortia enriched with different plant biomass reveal distinct players acting in lignocellulose degradation. *Microbial Ecology*, 71(3), 616–627. <https://doi.org/10.1007/s00248-015-0683-7>
- Dicko, M., Ferrari, R., Tangthirasunun, N., Gautier, V., Lalanne, C., Lamari, F., & Silar, P. (2020). Lignin degradation and its use in signaling development by the coprophilous ascomycete *Podospora anserina*. *Journal of Fungi*, 6(4), 278–301. <https://doi.org/10.3390/jof6040278>
- Gorea, E., Godwin, I., & Mudge, A. (2019). *Ganoderma* infection of oil palm – A persistent problem in Papua New Guinea and Solomon Islands. *Australasian Plant Pathology*, 49, 1–9. <https://doi.org/10.1007/s13313-019-00673-9>
- Hanis, A., Uke, A., Sudesh, K., & Kosugi, A. (2024). Accumulation of starch and sugars, and effect on pathogenesis-related proteins in felled oil palm trunks from the replanting period. *Industrial Crops and Products*, 218, 118863. <https://doi.org/10.1016/j.indcrop.2024.118863>
- Kang, J., Chu, Y., Ma, G., Zhang, Y., Zhang, X., Wang, M., Lu, H., Wang, L., Kang, G., Ma, D., Xie, Y., & Wang, C. (2023). Physiological mechanisms underlying reduced photosynthesis in wheat leaves grown in the field under conditions of nitrogen and water deficiency. *Crop Journal*, 1, 638–650. <https://doi.org/10.1016/j.cj.2022.06.010>

- Kuala Lumpur Kepong Berhad. (2018). *Sustainability report*. <http://myagric.upm.edu.my/id/eprint/20186/1/KLK-Sustainability-Report-2018a.pdf>
- Lai, J., Idris, A., & Yusof, N. M. (2014). Lignin extraction from oil palm trunk by microwave-alkali technique. *Malaysian Journal of Fundamental and Applied Sciences*, 10, 59. <https://doi.org/10.11113/mjfas.v10n2.56>
- Li, L., Xu, M., Ali, M., Zhang, W., Duan, Y., & Li, D. (2018). Factors affecting soil microbial biomass and functional diversity with the application of organic amendments in three contrasting cropland soils during a field experiment. *PLoS ONE*, 13, e0203812. <https://doi.org/10.1371/journal.pone.0203812>
- Lin, L., Xu, F., Ge, X., & Li, Y. (2019). Biological treatment of organic materials for energy and nutrients production – Anaerobic digestion and composting. In Y. Li & X. Ge (Eds.), *Advances in Bioenergy* (Vol. 4, pp. 121–181). Elsevier.
- Loh, S. K., James, S., Ngatiman, M., Cheong, K. Y., Choo, Y. M., & Lim, W. S. (2013). Enhancement of palm oil refinery waste – Spent bleaching earth (SBE) into bio organic fertiliser and their effects on crop biomass growth. *Industrial Crops and Products*, 49, 775–781. <https://doi.org/10.1016/j.indcrop.2013.06.016>
- Manjeri, G., Muhamad, R., & Tan, S. (2014). *Oryctes rhinoceros* beetles, an oil palm pest in Malaysia. *Annual Research Review of Biology*, 4(22), 3429–3439.
- McGuire, K., D'Angelo, H., Brearley, F., Gedallovich, S., Babar, N., Yang, N., Gillikin, C., Gradoville, M., Bateman, C., Turner, B., Mansor, P., Leff, J., & Fierer, N. (2014). Responses of soil fungi to logging and oil palm agriculture in Southeast Asian tropical forests. *Microbial Ecology*, 69(4), 733–747. <https://doi.org/10.1007/s00248-014-0468-4>
- Mohammed, A., Jamali, N., Yusop, M. A. S., & Adip, M. E. (2014). Effects of oil palm residues as bio-compost on growth performance of oil palm seedling. In *Prosiding Konferensi Akademik (KONAKA) 2014*, UiTM Pahang.
- Malaysian Palm Oil Board (2016). *MPOB Code of practice – Code of good nursery practice for oil palm nurseries* (2nd ed.).
- Pulingam, T., Lakshmanan, M., Chuah, J. A., Surendran, A., Zainab, L. I., Foroozandeh, P., Uke, A., Kosugi, A., & Sudesh, K. (2022). Oil palm trunk waste: Environmental impacts and management strategies. *Industrial Crops and Products*, 189, 115827. <https://doi.org/10.1016/j.indcrop.2022.115827>
- Qu, F., Peng, T., Jia, Y., Yang, M., Meng, X., Mao, S., Zhou, D., & Hu, X. (2022). Adjusting leaf nitrogen allocation could promote photosynthetic capacity, and nitrogen accumulation in *Cucumis sativus* L. *Environmental and Experimental Botany*, 198, 104855. <https://doi.org/10.1016/j.envexpbot.2022.104855>
- Ruan, Z., Zanotti, M., Wang, X., Ducey, C., & Liu, Y. (2012). Evaluation of lipid accumulation from lignocellulosic sugars by *Mortierella isabellina* for biodiesel production. *Bioresource Technology*, 110, 198–205. <https://doi.org/10.1016/j.biortech.2012.01.053>
- Savy, D., Mercl, F., Cozzolino, V., Spaccini, R., Cangemi, S., & Piccolo, A. (2020). Soil amendments with lignocellulosic residues of biorefinery processes affect soil organic matter accumulation and microbial growth. *ACS Sustainable Chemistry & Engineering*, 8(8), 3381–3391. <https://doi.org/10.1021/acssuschemeng.9b07474>
- Situmorang, E., Nugroho, Y., Prameswara, A., Andarini, E., Hartono, H., Hendroko, R., Toruan-Mathius, N., & Liwang, T. (2016). The bacterial diversity investigation in oil palm plantation using terminal restriction length polymorphism. In *Proceedings of the 4th International Conference on Biological Science*, (pp. 1–9). Universitas Gadjah Muda.
- Souza, C., Silva, T., de Souza, L., Moura, M., Pedro-Silva, P., & Marin, F. (2020). Straw management effects on sugarcane growth, nutrient cycling, and water use in the Brazilian semiarid region. *Bragantia*, 79(4), 400–411. <https://doi.org/10.1590/1678-4499.20200227>
- Sung, C. T. B. (2016). *Availability, use, and removal of oil palm biomass in Indonesia*. International Council on Clean Transportation. <https://doi.org/10.13140/RG.2.1.4697.4485>
- Uke, A., Nakazono-Nagaoka, E., Chuah, J. A., Zain, N. A. A., Amir, H. G., Sudesh, K., Abidin, N. Z. H. A. Z., Hashim, Z., & Kosugi, A. (2021). Effect of decomposing oil palm trunk fibres on plant growth and soil microbial community composition. *Journal of Environmental Management*, 295, 113050. <https://doi.org/10.1016/j.jenvman.2021.113050>

- Ungkulpasvich, U., Baramée, S., Uke, A., & Kosugi, A. (2021). Symbiotic chitin degradation by a novel anaerobic thermophilic bacterium *Hydrogenispora* sp. UUS1-1 and the bacterium *Tepidanaerobacter* sp. GT38. *Enzyme and Microbial Technology*, *144*, 109740. <https://doi.org/10.1016/j.enzmictec.2020.109740>
- Wen, B., Li, C., Fu, X., Li, D., Li, L., Chen, X., Wu, H., Cui, X., Zhang, X., Shen, H., Zhang, W., Xiao, W., & Gao, D. (2019). Effects of nitrate deficiency on nitrate assimilation and chlorophyll synthesis of detached apple leaves. *Plant Physiology and Biochemistry*, *142*, 363–371. <https://doi.org/10.1016/j.plaphy.2019.07.007>
- Wu, H., Cui, H., Fu, C., Li, R., Qi, F., Liu, Z., Yang, G., Xiao, K., & Qiao, M. (2024). Unveiling the crucial role of soil microorganisms in carbon cycling: A review. *Science of the Total Environment*, *909*, 168627. <https://doi.org/10.1016/j.scitotenv.2023.168627>
- Wuest, S., Albrecht, S., & Skirvin, K. (2000). Crop residue position and interference with wheat seedling development. *Soil and Tillage Research*, *55*(3-4), 175–182. [https://doi.org/10.1016/S0167-1987\(00\)00116-1](https://doi.org/10.1016/S0167-1987(00)00116-1)
- Yamada, H., Tanaka, R., Sulaiman, O., Hashim, R., Hamid, Z. A. A., Yahya, M. K. A., Kosugi, A., Arai, T., Murata, Y., Nirasawa, S., Yamamoto, K., Ohara, S., Mohd Yusof, M. N., Ibrahim, W. A., & Mori, Y. (2010). Old oil palm trunk: A promising source of sugars for bioethanol production. *Biomass and Bioenergy*, *34*(11), 1608–1613. <https://doi.org/10.1016/j.biombioe.2010.06.011>
- Zhang, Y., Wu, W., & Cai, L. (2017). Polyphasic characterisation of *Chaetomium* species from soil and compost revealed high number of undescribed species. *Fungal Biology*, *121*(1), 21–43. <https://doi.org/10.1016/j.funbio.2016.08.012>