

# SMALL SCALE FIELD EVALUATION OF A NOVEL PALM-BASED INSECTICIDE FORMULATION USING COLD AND THERMAL FOGGING AGAINST *Aedes aegypti*

SUMAIYAH MEGAT NABIL MOHSIN<sup>1\*</sup>; NORASHIKIN AHMAD<sup>1</sup> and YUSRABBIL AMIYATI YUSOF<sup>1</sup>

## ABSTRACT

*A new insecticide formulation with deltamethrin as an active ingredient (F1) has been developed in order to suppress the worldwide spread of mosquito-borne diseases. The formulation is naphtha-free and utilises locally sourced palm-based materials to obtain a cost-effective product. The objective of this study is to assess the performance of the formulation, in terms of droplet size and efficacy, in comparison to a commercial naphtha-based insecticide (F2). Open-field bioassay of the formulations against adult Aedes aegypti was conducted using cold and thermal fogging methods and measured at distance points 25, 50, 75 and 100 m. Overall, droplet size and efficacy significantly decreased with increasing distance from the spray path. No significant differences in volume median diameter (VMD) were observed between fogging methods and formulations. In general, both formulations showed significantly higher efficacy as cold fogs than thermal fogs. Significant differences in knockdown effect were observed between methods, time and formulations. Nevertheless, the definite effect of insecticidal formulations, i.e. mortality, was not significantly different between F1 and F2. In conclusion, our results show that the palm-based formulation was able to achieve comparable performance with the commercial formulation without the use of naphtha.*

**Keywords:** deltamethrin, droplet size, efficacy, space spray, vector control.

**Received:** 24 July 2024; **Accepted:** 13 January 2025; **Published online:** 19 March 2025.

## INTRODUCTION

Dengue, an arboviral disease, has spread rapidly across the globe and is becoming endemic in more than 100 countries (Selvarajoo et al., 2020). Regular dengue outbreaks have occurred in Malaysia since the 1960s, and the case numbers continue to increase alongside Malaysia's rapid population and infrastructure growth. Malaysia recorded the biggest spike in dengue cases in four years in 2019 (AbuBakar et al., 2022). Dengue vector control activities in Malaysia rely heavily on human resources (Yazan et al., 2021). Thus, lockdowns or movement control order (MCO) to contain the spread of Coronavirus disease 2019 (COVID-19) have significantly curtailed regular dengue control

operations (Rahim et al., 2021). At present, there are no effective treatments available for dengue. Furthermore, the current vaccine exhibits moderate efficacy and fails to provide balanced protection against all four dengue serotypes (Stanaway et al., 2016). Hence, control of the dengue vector population remains the mainstay strategy for containing the spread of the disease.

Space spraying using thermal and cold foggers is an often-used intervention strategy for controlling mosquito-borne diseases (Pryce et al., 2018). However, space sprays are only effective when the droplets remain airborne. Thus, multiple applications are necessary for effective control of the dengue vector population (World Health Organization [WHO], 2009). With the ever-increasing fuel cost and concerns for adverse environmental impacts from frequent space spray interventions, the use of water-dilutable water-based formulation is turning into an appealing and trendy alternative to oil-

<sup>1</sup> Malaysian Palm Oil Board,  
6, Persiaran Institusi, Bandar Baru Bangi,  
43000 Kajang, Selangor, Malaysia.

\* Corresponding author e-mail: [sumaiyah@mpob.gov.my](mailto:sumaiyah@mpob.gov.my)

based formulation. Questions have been raised about the efficacy of water-based formulations as the vapour pressure of water causes droplet evaporation and consequently size reduction, creating insecticide droplets that are prone to drift (Farajollahi & Williams, 2013). Nevertheless, efficacy studies using water-based formulations have demonstrated that they are equally or more effective than formulations using diesel fuel or mineral oil as a diluent (Farajollahi & Williams, 2013; Harburguer et al., 2012a).

In recent years, bio-based formulations incorporating plant and microbial bioactives have been explored to address environmental impacts and insecticide resistance in mosquito control. Bioformulations of plant extracts from *Carica papaya* and *Parthenium hysterophorus* have shown promising larvicidal activity against *Aedes aegypti* (Jayaraman et al., 2023). Entomopathogenic fungi, *Beauveria bassiana* JN5R1W1, when formulated as inverted emulsions with soybean oil demonstrate high mortality rates against adult mosquitoes through both direct and indirect contact methods (Yong-Lee et al., 2023). Under the field-simulated trial, *Ocimum kilimandscharicum* oil formulation achieved 98% larvicidal mortality against *Anopheles gambiae* larvae after 24 hr, as opposed to 54% mortality achieved by *Bacillus thuringiensis* subsp. *israelensis* granules (Ochola et al., 2022). While these bioformulations comprising bioactives show promise, they often lack the robustness and field applicability of synthetic chemicals. Thus, while bioactives provide potential supplemental strategies, synthetic chemicals remain the cornerstone of effective vector control in the field.

In response to these challenges, a novel water-based insecticidal formulation (F1) comprising palm-based solvents and surfactants has been developed to restrain dengue outbreaks and minimise the environmental impact of vector control (Mohsin et al., 2019). The formulation contains deltamethrin, a larvicide and adulticide that causes an instant knockdown by paralysing the nervous system of insects (Kumar et al., 2011). F1 is free from highly volatile and flammable naphtha solvent and incorporates palm-based methyl esters (PME) as a substitute. PME is less volatile and possesses higher flash point compared to petroleum-based solvents (Mohsin et al., 2017). Additionally, F1 comprises non-toxic and readily biodegradable palm-based surfactants which are instrumental in improving the stability of pesticide emulsions (Ghazali & Ahmad, 2004; Ismail et al., 2014; Shaari et al., 2022; Siti Afida et al., 2016). Malaysia's export revenue of palm-based oleochemicals in 2023 declined by 30.1% from the previous year despite the marginal improvement in export volume (4.8%) due to lower imports by the major buyers of Malaysian palm oil

(Parveez et al., 2024). Given the recent decline, leveraging these locally sourced bio-based components offers both economic and strategic benefits by reducing dependency on imported hydrocarbons.

This study addresses the novel application of F1 as an environmentally friendly and cost-effective solution for dengue vector control. By comparing F1 to a commercially available deltamethrin formulation (F2) in an open-field setting, we aim to evaluate their respective efficacies in terms of droplet size distribution, knockdown and mortality of the primary dengue virus vector *Ae. aegypti* using cold and thermal fogging methods. This investigation is critical in determining whether a palm-based water-dilutable insecticide can meet or exceed the performance of conventional formulations, providing a new, sustainable avenue for managing dengue outbreaks in Malaysia and other endemic regions.

## MATERIALS AND METHODS

### Biological Material

Sucrose-fed adult *Ae. aegypti* females (aged 2–5 days) from the Vector Control Research Unit, Universiti Sains Malaysia (USM), Penang, Malaysia, were used for the adulticidal assessments.

### Chemicals

PME comprising a mixture of methyl caproate, methyl octanoate, methyl decanoate and methyl laurate at 3.2 wt%, 50.8 wt%, 44.0 wt% and 2.0 wt%, respectively, was provided by Emery Oleochemicals Sdn. Bhd., Malaysia. Deltamethrin (98.0%) was donated by Hextar Chemicals Sdn. Bhd. Malaysia. Xanthan gum derived from *Xanthomonas campestris* was purchased from Sigma-Aldrich Sdn. Bhd., Malaysia. Bio-based surfactants, C12–C14 fatty alcohol ethoxylate and polyoxyethylene (20) sorbitan monooleate, are used in the study. The former was obtained from Huntsman Singapore Pte. Ltd., Singapore, while the latter was purchased from Chemweb Sdn. Bhd., Malaysia.

### Insecticide Formulations

F1 was prepared using the method in a previous study by Mohsin et al., (2019). In brief, the formulation was developed using 2.0 wt% deltamethrin, 5.0 wt% mixture of the previously mentioned bio-based surfactants and 13.0 wt% PME as the oil phase and distilled water and 0.4 wt% xanthan gum as the aqueous phase. The hydrophilic-lipophilic balance of the surfactant

was kept constant at 10. The oil phase was prepared by stirring for 10 min at 55°C, while the aqueous phase was prepared by dispersing xanthan gum in distilled water at 55°C. Thereon, the oil phase was gradually incorporated into the aqueous phase using a Kinematica Polytron® PT3100 homogeniser (Kinematica AG, Switzerland) operated at 7,000 rpm for 10 min. The commercial insecticide (F2) is an oil-in-water (o/w) emulsion formulation containing 2.0 wt% deltamethrin and petroleum-based naphtha from Bayer Environmental Science, Spain. The blank formulation (F3) was similarly produced according to the previously mentioned process without deltamethrin to examine the possible insecticidal activity of the emulsion system.

### Fogging Method

All insecticide formulations were water-diluted at a dilution ratio of 1:100 v/v. Then, thermal fogging was applied at 5,000 mL ha<sup>-1</sup> using a hand carried Agrofog Hand Held Thermal Fogger (Agro Technic Pte. Ltd., Singapore). On the other hand, cold fogging was performed at 500 mL ha<sup>-1</sup> using a truck mounted LECO ULV Fog Generator Model 1800 (Clarke, USA) with its sprayer head nozzle angled 30° upwards to the horizontal plane. The truck then travelled 6–9 km hr<sup>-1</sup> at a distance of 200 m perpendicular to the spray angle.

### Droplet Size Measurement and Efficacy Assessment

Field tests were performed in accordance with the WHO guidelines WHO (2009) in an open field in USM, with no tall grasses that should hinder fog movement. The study was conducted during fair weather. The temperature during treatment was 27°C–30°C, the relative humidity was 70%–80% and the wind speed was 1.5–3.0 m s<sup>-1</sup>. A total of 20 mosquitoes were held in nylon mesh-screened cylindrical cages (1.2 mm mesh size) with wire frame support (10 cm diameter × 15 cm height × 10 cm tapping cover). The cages were suspended at 1.5 m from the ground on poles distanced at 25, 50, 75 and 100 m from the spray nozzle. Knockdown was recorded after 15 min, and the mosquitoes were transferred into mesh covered cups using battery operated aspirator and returned to the laboratory. The cups were provided with a cotton pad soaked with 10% sucrose solution. Then, knockdown was recorded again at 30 min and 60 min post-treatment while adult mortality was recorded at 24 hr post-treatment. Three replicates were conducted for each formulation (including water control i.e., water without insecticide). Results from the water control treatment were excluded from reporting as knockdown and mortality activities were not observed. In each test, one set of adult cages was

kept in the laboratory as a control. If mortality was above 20% in the concurrent controls, the whole test would be rejected. Results from the treated samples will be corrected using Abbott's formula if mortality in the controls was above 5% (Abbott, 1925). Abbott's formula for calculating corrected mortality is as Equation (1):

$$\frac{\% \text{ Mortality in treated} - \% \text{ Mortality in control}}{100\% - \% \text{ Mortality in control}} \times 100 \quad (1)$$

To record fog droplet sizes at varied pole distances, magnesium oxide (MgO)-coated slides were placed approximately 1.5 m above the ground on a slide rotator at each pole. The droplets were examined under a microscope and their volume medium diameter (VMD) in mm was calculated using Microsoft Excel developed by Clarke Engineering Technologies, USA.

### Statistical Analysis

Data were analysed using IBM SPSS Statistics 25 (IBM, USA). Two-way analysis of variance (ANOVA) followed by Tukey's Honestly-Significant-Difference (Tukey HSD) post-hoc test was carried out to determine significant differences in fog droplet sizes between formulations (F1, F2 and F3) and distance (25, 50, 75 and 100 m). Similarly, two-way ANOVA and Tukey HSD were carried out to determine significant differences in the efficacy results (knockdown and mortality) between distance and knockdown time (15, 30 and 60 min). Differences between means were considered statistically significant at  $p < 0.05$ . In addition, t-tests were carried out to compare the means of droplet sizes between fogging methods and means of efficacy results between formulations (F1 and F2) and fogging methods.

## RESULTS AND DISCUSSION

### Droplet Size

Fog droplet size is an important factor influencing droplet deposition and the efficacy of insecticides. During space spraying, large droplets may fail to stay airborne or penetrate obstacles, while very small droplets may fail to land on target insects due to aerodynamics and convection currents (Farajollahi & Williams, 2013). VMD is a commonly used parameter to describe the droplet diameter, i.e., where 50% of the spray liquid volume consists of droplets smaller than this value (He et al., 2022). Accordingly, the optimum droplet size for efficacious ground adulticiding was

determined to be between 12 and 20  $\mu\text{m}$  VMD (Bonds, 2012). The VMD of droplets produced in this study are within said range until up to 50 m but decline below the optimal size at further distances (Table 1).

Two-way ANOVA analysis showed no significant difference in VMD between formulations (Table 2), indicating that droplets produced using palm-based formulations F1 and F3 were comparable to the commercial product, F2. The water-based formulations exhibited significant droplet size reduction with increasing distance. The finding is in line with previous studies by Bengoa et al. (2014) and Farajollahi and Williams (2013), which attributed the droplet size reduction to water evaporation in the outdoor environment as they travelled further away from the spray nozzle. Additionally, t-test analysis revealed that there is no significant difference in the droplet sizes produced by cold and thermal foggers (Table 2), which is in agreement with the findings obtained by Harburguer et al. (2012b) and Megat Nabil Mohsin et al. (2023). There was a statistically significant interaction between method and distance. In contrast, the interaction between formulation and distance and the interaction between formulation and method were non-significant (Table 2).

### Knockdown of Adult Mosquitoes

Quick knockdown from exposure to insecticidal formulation ensures a reduction in feeding and insect reproduction and diminishes the chances for insects to escape (Georgia et al., 2018). In the

absence of deltamethrin, F3 showed negligible knockdown with the maximum being only 3.33% at all checkpoints via cold fogger and 1.67% at the 25 m checkpoint via thermal fogger after 60 min (Figure 1). The results showed that the palm-based emulsion system alone had no knockdown activity and thus is excluded from statistical analysis.

A significant decrease in knockdown was observed as the distance increased, similar to the trend reported for droplet size. However, unlike droplet size, knockdown was significantly different between methods and between formulations F1 and F2 (Table 3). With cold fogging, knockdown by F1 was slightly lower than the commercial insecticide at the 15 min interval (Figure 1a). However, the opposite result was observed with thermal fogging (Figure 1b). In general, it was observed that knockdown significantly increased with time. At the 30 and 60 min intervals, cold fogging using F1 and F2 had achieved 100% knockdown at all checkpoints intervals, indicative of a low recovery rate for the knocked-down mosquitoes. Nevertheless, lower knockdown percentages were observed with thermal fogging compared to cold fogging as both F1 and F2 had failed to achieve complete knockdown even at 60 min (Figure 1b).

Two-way ANOVA revealed significant interactions between method and distance, method and formulation, method and time, time and distance, as well as time and formulation. However, there was no significant difference in the interaction between formulation and distance (Table 3). A note of caution is due here since the knockdown effect, i.e., the paralysis of insects due to insecticides

TABLE 1. DROPLET MEASUREMENT AT SPECIFIED DISTANCES USING DIFFERENT FOGGING METHODS

Fogging method	Formulation	VMD ( $\mu\text{m}$ ) (mean $\pm$ S.E.)			
		25 m	50 m	75 m	100 m
Cold	F1	13.40 $\pm$ 0.12	12.60 $\pm$ 0.10	11.57 $\pm$ 0.06	10.20 $\pm$ 0.06
	F2	13.40 $\pm$ 0.06	12.73 $\pm$ 0.03	11.50 $\pm$ 0.06	10.23 $\pm$ 0.09
	F3	13.33 $\pm$ 0.12	12.63 $\pm$ 0.03	11.97 $\pm$ 0.18	10.10 $\pm$ 0.00
Thermal	F1	12.83 $\pm$ 0.12	12.30 $\pm$ 0.12	11.87 $\pm$ 0.20	10.30 $\pm$ 0.06
	F2	12.70 $\pm$ 0.10	12.27 $\pm$ 0.03	12.07 $\pm$ 0.09	10.30 $\pm$ 0.12
	F3	12.87 $\pm$ 0.07	12.23 $\pm$ 0.07	11.90 $\pm$ 0.06	10.30 $\pm$ 0.06

Note: F1 - novel water-based insecticidal formulation; F2 - commercial insecticide; F3 - blank formulation; VMD - volume median diameter; S.E. - standard error.

TABLE 2. THE EFFECTS OF FOGGING METHOD, FORMULATION AND DISTANCE ON FOG DROPLET SIZE

Variable	df	F-value	P-value
Fogging method	70	4.417	0.580 (2-tailed)
Formulation	2	0.219	0.804
Distance	3	885.136	<0.001
Fogging method $\times$ distance	3	24.004	<0.001
Fogging method $\times$ formulation	2	0.237	0.790
Formulation $\times$ distance	6	0.972	0.453

Note: df - degree of freedom.


is potentially reversible (Tsaganou et al., 2021). Therefore, we must also look into the delayed but definite effect of insecticidal formulations, i.e., mortality.

### Adult Mortality

The objective of chemical space spraying is to minimise disease transmission by reducing the target infectious adult mosquitoes and their populations (Zuharah et al., 2019). As shown in Figure 2, negligible adult mortality was observed

with F3, ranging from 0%–3% for cold fogging and 0%–5% for thermal fogging. This result further reinforces that the palm-based emulsion system used in the formulation provided no insecticidal activity.

The statistical analysis demonstrated no significant differences in adult mortality between F1 and F2 (Table 4). Despite the absence of highly volatile and flammable naphtha solvents, the palm-based formulation was able to provide comparable adulticidal efficacy with the commercial formulation. In this study, mortality



Note: F1 - novel water-based insecticidal formulation; F2 - commercial insecticide; F3 - blank formulation.

Figure 1. Percentage knockdown (mean ± S.E.) of adult *Ae. aegypti* after exposure to fogging formulations using (a) cold and (b) thermal fogger.

TABLE 3. THE EFFECTS OF FOGGING METHOD, FORMULATION, KNOCKDOWN TIME AND DISTANCE ON PERCENTAGE KNOCKDOWN

Variable	df	F-value	P-value
Fogging method	142	97.787	<0.001 (2-tailed)
Formulation	142	16.819	0.049 (2-tailed)
Knockdown time	2	24.530	<0.001
Distance	3	60.943	<0.001
Fogging method × distance	3	44.297	<0.001
Fogging method × formulation	1	36.872	<0.001
Fogging method × time	2	7.390	0.001
Knockdown time × distance	6	2.378	0.033
Knockdown time × formulation	2	4.279	0.016
Distance × formulation	3	2.500	0.063


Note: df - degree of freedom.

was seen to decline significantly with increasing distance, a similar pattern was observed with knockdown. This may be attributed to droplet size reduction with increasing travel distance. Larger droplets are able to accomplish higher efficacies compared to smaller droplets because they are harder to transport by air currents because of their larger inertial force (Sugiura et al., 2011). In contrast to knockdown, mortality was significantly different between fogging methods (Table 4). It is worth highlighting that the cold fogging method was more effective in controlling adult *Ae. aegypti*, achieving 100% mortalities at all distance points (Figure 2). This pattern is in accord with previous studies, which attributed the poorer performance of their water-based products via thermal fogging to excessive heat by the applicator, resulting in reduced efficacy (Fulcher et al., 2016; Megat Nabil Mohsin et al., 2023). Nevertheless, additional studies are required to assess the efficacies of this palm-based formulation against immature *Ae. aegypti* stages, pupae and larvae.

Two-way ANOVA revealed that similarly with knockdown, a significant interaction was observed between method and distance, whereas no significant interaction was observed between formulation and distance. Nonetheless, contrary to the trends observed with knockdown, a significant difference was not observed in the interaction between method and formulation (Table 4).

## CONCLUSION

In conclusion, this study successfully demonstrated that the novel palm-based insecticide formulation (F1) offers comparable efficacy to the commercial naphtha-based formulation (F2) in controlling *Ae. aegypti* mosquitoes, without the environmental and safety concerns associated with petroleum-derived solvents. The field trials using both cold and thermal fogging methods showed no significant differences in droplet sizes between formulations, while the performance of cold fogging was generally superior in terms of knockdown and mortality. Importantly, this research highlights the potential of utilising palm-based ingredients, particularly PME and surfactants, which are biodegradable and locally abundant, reducing dependence on non-renewable resources. The use of F1 also provides a cost-effective and sustainable alternative for dengue vector control, aligning with global trends toward greener, more environmentally responsible pest control solutions. This study is unique in its focus on replacing volatile, flammable naphtha with a bio-based formulation, which is particularly relevant in regions like Malaysia, where palm oil is a major agricultural product. The findings suggest that palm-based insecticidal formulations could play a significant role in future vector control strategies, particularly in tropical and subtropical regions where mosquito-borne diseases are prevalent.



Note: F1 - novel water-based insecticidal formulation; F2 - commercial insecticide; F3 - blank formulation.

Figure 2. Percentage mortality (mean  $\pm$  S.E.) of adult *Ae. aegypti* after exposure to fogging formulations using cold and thermal foggers.

TABLE 4. THE EFFECTS OF FOGGING METHOD, FORMULATION AND DISTANCE ON ADULT MORTALITY

Variable	df	F-value	P-value
Fogging method	46	57.982	0.004 (2-tailed)
Formulation	46	0.239	0.725 (2-tailed)
Distance	3	27.445	<0.001
Fogging method $\times$ distance	3	27.445	<0.001
Fogging method $\times$ formulation	1	0.665	0.420
Distance $\times$ formulation	3	0.955	0.425

Note: df - degree of freedom.

## ACKNOWLEDGEMENT

The authors wish to express their gratitude to the Director-General of MPOB for granting permission to publish this article. The authors are grateful to Dr. Suhaida Abdullah, School of Quantitative Sciences, Universiti Utara Malaysia, for providing statistical consultations. The mention of a commercial product in this study does not constitute an endorsement. The authors have no financial interest in this product, and ties to this product are research in nature only.

## REFERENCES

- Abbott, W. S. (1925). A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*, 18(2), 265–267. <https://doi.org/10.1093/jee/18.2.265a>
- AbuBakar, S., Puteh, S. E. W., Kastner, R., Oliver, L., Lim, S. H., Hanley, R., & Gallagher, E. (2022). Epidemiology (2012–2019) and costs (2009–2019) of dengue in Malaysia: A systematic literature review. *International Journal of Infectious Diseases*, 124, 240–247. <https://doi.org/10.1016/j.ijid.2022.09.006>
- Bengoa, M., Eritja, R., & Lucientes, J. (2014). Ground ultra-low volume adulticiding field trials using pyrethroids against *Aedes albopictus* in the Baix Llobregat Region, Spain. *Journal of the American Mosquito Control Association*, 30(1), 42–50. <https://doi.org/10.2987/13-6378.1>
- Bonds, J. A. S. (2012). Ultra-low-volume space sprays in mosquito control: A critical review. *Medical and Veterinary Entomology*, 26(2), 121–130. <https://doi.org/10.1111/j.1365-2915.2011.00992.x>
- Farajollahi, A., & Williams, G. M. (2013). An open-field efficacy trial using AquaDuet™ via an ultra-low volume cold aerosol sprayer against caged *Aedes albopictus*. *Journal of the American Mosquito Control Association*, 29(3), 304–308. <https://doi.org/10.2987/13-6334R.1>
- Fulcher, A., Farooq, M., Richardson, A. G., Smith, M. L., Scott, J. M., Gaines, M. K., & Xue, R. D. (2016). Characteristics and efficacy of three commercial handheld thermal foggers with pyrethroid insecticides against three species of mosquitoes. *Journal of the American Mosquito Control Association*, 32(1), 44–50. <https://doi.org/10.2987/8756-971X-32.1.44>
- Georgia, B., Christos, R. I., & Christos, A. G. (2018). From lethality to mortality: Exploring the “grey area” of knockdown as an efficacy indicator of different insecticides against major storage insects using a lethality index. *Journal of Pest Science*, 91(4), 1371–1380. <https://doi.org/10.1007/s10340-018-0983-z>
- Ghazali, R., & Ahmad, S. (2004). Biodegradability and ecotoxicity of palm stearin-based methyl ester sulfonates. *Journal of Oil Palm Research*, 16(1), 39–44.
- Harburguer, L., Lucia, A., Licastro, S., Zerba, E., & Masuh, H. (2012a). Field comparison of thermal and non-thermal ultra-low-volume applications using water and diesel as solvents for managing dengue vector, *Aedes aegypti*. *Tropical Medicine & International Health*, 17(10), 1274–1280. <https://doi.org/10.1111/j.1365-3156.2012.03059.x>
- Harburguer, L., Seccacini, E., Licastro, S., Zerba, E., & Masuh, H. (2012b). Droplet size and efficacy of an adulticide-larvicide ultralow-volume formulation on *Aedes aegypti* using different solvents and spray application methods. *Pest Management Science*, 68(1), 137–141. <https://doi.org/10.1002/ps.2237>
- He, Y., Wu, J., Fu, H., Sun, Z., Fang, H., & Wang, W. (2022). Quantitative analysis of droplet size distribution in plant protection spray based on machine learning method. *Water*, 14(2), 175. <https://doi.org/10.3390/w14020175>
- Ismail, A. R., Nor Farhana, N., Mahiran, B., Dzolkhifli, O., & Hazimah, A. H. (2014). Oil-in-water emulsion (EW) of mixed glyphosate isopropylamine (IPA) and triclopyr butoxyethylester (BEE) stabilized by palm-based emulsifiers for weed control. *Journal of Oil Palm Research*, 26(4), 366–374.
- Jayaraman, T., Chandrapragasam, V., & Kavitha, S. (2023). Development of bioformulations using plant extracts for the control of dengue vector, *Aedes aegypti*. *Journal of Applied and Natural Science*, 15(2), 760–766. <https://doi.org/10.31018/jans.v15i2.4518>
- Kumar, S., Thomas, A., & Pillai, M. K. K. (2011). Deltamethrin: Promising mosquito control agent against adult stage of *Aedes aegypti* L. *Asian Pacific Journal of Tropical Medicine*, 4(6), 430–435. [https://doi.org/10.1016/S1995-7645\(11\)60120-X](https://doi.org/10.1016/S1995-7645(11)60120-X)
- Megat Nabil Mohsin, S., Ahmad, N., & Yusof, Y. A. (2023). Comparative efficacy trials with alcohol added d-phenothrin formulations against *Aedes aegypti* under open-field condition. *Pest Management Science*, 79(10), 4094–4099. <https://doi.org/10.1002/ps.7610>

- Mohsin, S. M. N., Raman, I. A., Hasan, Z. A. A., & Idris, Z. (2017). Palm-based methyl esters as carrier solvents in pesticide formulations. *Palm Oil Developments*, 66, 32–38.
- Mohsin, S. M. N., Raman, I. A., Hasan, Z. A. A., & Idris, Z. (2019). Influence of surfactant structure on physical stability and physicochemical properties of insecticide emulsions. *Journal of Oil Palm Research*, 31(4), 670–680. <https://doi.org/10.21894/jopr.2019.0025>
- Ochola, J. B., Mutero, C. M., Marubu, R. M., Haller, B. F., Hassanali, A., & Lwande, W. (2022). Mosquitoes larvicidal activity of *Ocimum kilimandscharicum* oil formulation under laboratory and field-simulated conditions. *Insects*, 13(2), 203. <https://doi.org/10.3390/insects13020203>
- Parveez, G. K. A., Leow, S.-S., Kamil, N. N., Madihah, A. Z., Ithnin, M., Ng, M. H., Yusof, Y. A., & Idris, Z. (2024). Oil palm economic performance in Malaysia and R&D progress in 2023. *Journal of Oil Palm Research*, 36(2), 171–186. <https://doi.org/10.21894/jopr.2024.0037>
- Pryce, J., Choi, L., Richardson, M., & Malone, D. (2018). Insecticide space spraying for preventing malaria transmission. *Cochrane Database of Systematic Reviews*, 2018(11), 1–46. <https://doi.org/10.1002/14651858.CD012689.pub2>
- Rahim, M. H., Dom, N. C., Ismail, S. N. S., Mulud, Z. A., Abdullah, S., & Pradhan, B. (2021). The impact of novel coronavirus (2019-nCoV) pandemic movement control order (MCO) on dengue cases in Peninsular Malaysia. *One Health*, 12, 100222. <https://doi.org/10.1016/j.onehlt.2021.100222>
- Selvarajoo, S., Liew, J. W. K., Tan, W., Lim, X. Y., Refai, W. F., Zaki, R. A., Sethi, N., Wan Sulaiman, W. Y., Lim, Y. A. L., Vadivelu, J., & Vythilingam, I. (2020). Knowledge, attitude and practice on dengue prevention and dengue seroprevalence in a dengue hotspot in Malaysia: A cross-sectional study. *Scientific Reports*, 10, 9534. <https://doi.org/10.1038/s41598-020-66212-5>
- Shaari, A. L., Zolkarnain, N., Raman, I. A., & Ghazali, R. (2022). Effects of palm oil-based surfactant used in pesticide formulation on terrestrial plants. *Journal of Oil Palm Research*, 34(4), 721–730. <https://doi.org/10.21894/jopr.2022.0017>
- Siti Afida, I., Razmah, G., & Zulina, A. M. (2016). Biodegradation of various homologues of palm-based methyl ester sulphonates (MES). *Sains Malaysiana*, 45(6), 949–954.
- Stanaway, J. D., Shepard, D. S., Undurraga, E. A., Halasa, Y. A., Coffeng, L. E., Brady, O. J., Hay, S. I., Bedi, N., Bensenor, I. M., Castañeda-Orjuela, C. A., Orjuela, C. A., Chuang, T. W., Gibney, K. B., Memish, Z. A., Rafay, A., Ukwaja, K. N., Yonemoto, N., & Murray C. J. L. (2016). The global burden of dengue: An analysis from the global burden of disease study 2013. *Lancet Infectious Diseases*, 16(6), 712–723. [https://doi.org/10.1016/S1473-3099\(16\)00026-8](https://doi.org/10.1016/S1473-3099(16)00026-8)
- Sugiura, M., Horibe, Y., Kawada, H., & Takagi, M. (2011). Effect of different droplet size on the knockdown efficacy of directly sprayed insecticides. *Pest Management Science*, 67(9), 1115–1123. <https://doi.org/10.1002/ps.2157>
- Tsaganou, F. K., Vassilakos, T. N., & Athanassiou, C. G. (2021). Knockdown-mortality responses of six stored-product beetle species on different surfaces treated with thiamethoxam, using a standardized lethality index. *Journal of Economic Entomology*, 114(1), 430–439. <https://doi.org/10.1093/jee/toaa272>
- World Health Organization. (2009). *Guidelines for efficacy testing of insecticides for indoor and outdoor ground-applied space spray applications*. [http://iris.who.int/bitstream/handle/10665/70070/WHO\\_HTM\\_NT-D\\_WHOPEPES\\_2009.2\\_end.pdf?sequence=1](http://iris.who.int/bitstream/handle/10665/70070/WHO_HTM_NT-D_WHOPEPES_2009.2_end.pdf?sequence=1)
- Yazan, L. S., Gopalsamy, B., Bakar, S. N. A., Manan, K. A., Shahidan, N. S., & Wang, L. Y. (2021). Efficacy of intelligent mosquito system (I.M.O.S) with XMOS mini aerosol against *Aedes* in 17th College, Universiti Putra Malaysia. *Pertanika Journal of Science and Technology*, 29(2), 1229–1242. <https://doi.org/10.47836/pjst.29.2.29>
- Yong-Lee, J., Mi Woo, R., & Dong Woo, S. (2023). Formulation of the entomopathogenic fungus *Beauveria bassiana* JN5R1W1 for the control of mosquito adults and evaluation of its novel applicability. *Journal of Asia Pacific Entomology*, 26(2), 102056. <https://doi.org/10.1016/j.aspen.2023.102056>
- Zuharah, W. F., Sumayyah, A., & Dieng, H. (2019). Susceptibility and fitness cost of *Aedes albopictus* on their survivability after exposure to the insecticide. *Journal of Asia Pacific Entomology*, 22(2), 553–560. <https://doi.org/10.1016/j.aspen.2019.03.003>