

CYTOTOXIC AND ANTI-ACNE ACTIVITY OF SUGAR FATTY ACID ESTERS (SFAE) FROM CPO AND PALM OIL HYDROLYSATE

TUTI WUKIRSARI^{1*}; ADE ANDRIANI²; OBIE FAROBIE³; SRI HANDAYANI¹ and SUMI HUDIYONO¹

ABSTRACT

This study aimed to determine the potential of sugar fatty acid esters (SFAE) derived from crude palm oil (CPO) and palm oil hydrolysates as cytotoxic and anti-acne agents. The influence of structural variations among different saccharides on the SFAE properties was also investigated. Three types of saccharides were tested in this study; D-mannitol (a sugar alcohol), D-mannose (an aldose) and D-fructose (a ketose). SFAEs were prepared enzymatically by reacting fatty acids and saccharides using the low-cost liquid lipase Eversa[®] Transform 2.0 at 5% (w/w) concentration. Thin layer chromatography (TLC) analysis, Fourier transform infrared spectroscopy (FTIR) and nuclear magnetic resonance (NMR) spectra, emulsification properties, conversion rate and degree of esterification indicated the formation of SFAE. Some SFAE exhibited strong cytotoxic and antibacterial activities. SFAE of palmitic acid-mannitol and SFAE of stearic acid-fructose showed the highest cytotoxic activity against MCF-7 cells, with an IC₅₀ value of 29 ppm. The SFAE of palm oil hydrolysate exhibited stronger cytotoxic activity than the SFAE of CPO hydrolysate. Additionally, the SFAE from CPO hydrolysate and D-mannitol exhibited the strongest antibacterial activity, achieving 100% inhibition against Cutibacterium acnes at 0.9 mg/mL. These findings highlight the potential of SFAE as a promising raw materials for the development of health and cosmetic products.

Keywords: CPO and palm oil hydrolysate, cytotoxic and anti-acne activities, enzymatic esterification, Eversa[®] Transform 2.0, sugar fatty acid ester.

Received: 25 July 2024; **Accepted:** 21 April 2025; **Published online:** 16 July 2025.

INTRODUCTION

Indonesia and Malaysia are the two largest palm oil producers in the world, accounting for most of global production (U.S. Department of Agriculture, 2024). These two countries dominate the market because of their favourable climates for palm oil cultivation and large-scale plantations. However, by the end of

2022, the European Union issued a new regulation on deforestation-free products that tightened the rules for importing crude palm oil (CPO) into its territory. Therefore, downstreaming palm oil and CPO into derivative products (oleochemicals) is crucial in increasing their sales value and benefits. The most common oleochemical products were fatty acid hydrolysates, glycerin, soap noodles and oleic acid. Oleochemical products in the form of sugar fatty acid esters (SFAE) have not been widely studied. One of the companies operating in this field is Ryoto[™] Sugar Ester, which is based in Japan. The main application of sucrose-fatty acid esters from Ryoto[™] is as an emulsifier in the food sector.

The application of SFAE as an emulsifier in the food industry showed that these compounds are safe for consumption (Sardar et al., 2024). The emulsification properties of SFAE have also been applied to the pharmaceutical, detergent and cosmetics industries (Qi et al., 2024). The non-ionic surfactant SFAE has also been used to produce niosomes for quercetin encapsulation (Elmowafy

¹ Department of Chemistry,
Faculty of Mathematics and Sciences,
Universitas Indonesia, Depok,
West Java 16424, Indonesia.

² Research Centre for Biotechnology,
National Research and Innovation Agency,
Jl. Raya Bogor, Cibinong, Bogor,
West Java 16911, Indonesia.

³ Department of Mechanical and Biosystem Engineering,
Faculty of Agricultural Engineering and Technology,
IPB University (Bogor Agricultural University),
IPB Dramaga Campus, Bogor,
West Java 16680, Indonesia.

* Corresponding author e-mail: tutiwukirsari@sci.ui.ac.id

et al., 2020). Niosome-encapsulated quercetin showed a higher hepatoprotective activity than free quercetin. Emulsification properties have also been reported to be associated with biological activities, such as antibacterial (Falk, 2019; Pirog et al., 2019) and anticancer (Chiewpattanakul et al., 2010; Walvekar et al., 2022). Emulsifiers or surfactants can damage the membranes of both microbes and cancer cells. The cytotoxic activity of SFAE has been described by Teng et al. (2021), Snoch et al. (2021), Chen et al. (2023) and Wang et al. (2024), while the antimicrobial activity of SFAE has been reported by Matin et al. (2020, 2022), Teng et al. (2021), and Yang et al. (2024). The role of SFAE in suppressing inflammation caused by acne bacteria was even reported by a US patent in 2005 (Pauly & Grisoni, 2005). However, study on the bioactivity comparison of SFAE from CPO hydrolysate and palm oil hydrolysate for anticancer and anti-acne has never been reported. Study into developing SFAE as a safe active compound is expected to be an alternative to anticancer and anti-acne medication. Anticancer and anti-acne medicines with minimal side effects are urgently needed because recent research by Vitiello et al. (2024) even shows that anticancer drugs can change the ocular surface of cancer chemotherapy patients. Wiśniewska et al. (2023) mentioned that salicylic acid, a well-known anti-acne cosmetic ingredient, causes several side effects such as hyperpigmentation, allergic reaction and swelling.

SFAE can be prepared either chemically or enzymatically. The chemical synthesis of SFAE has been described using a base-catalysed transesterification reaction between triglycerides or fatty acid methyl esters and sugars (Teng et al., 2021). In contrast, lipases have been widely reported for the enzymatic synthesis of SFAE. Surprisingly, proteases also exhibited the ability to catalyse the formation of SFAE (Bernal et al., 2018). Commercially available lipases that are widely used for the synthesis of SFAE include immobilised lipase B derived from *Candida antarctica* (Chen et al., 2023; Hollenbach et al., 2020; Nieto et al., 2021; Shin et al., 2019; Snoch et al., 2021) and immobilised lipase derived from *C. rugosa* (El-Baz et al., 2021). Ferdjani and Selaimia (2024) prepared SFAE using lipase from maize seed. The utilisation of lipase Eversa® Transform 2.0 as a catalyst for synthesising SFAE has never been studied. Eversa® Transform 2.0 is a low-cost liquid lipase for biodiesel production (Chang et al., 2021; Farobie et al., 2021). Monteiro et al. (2024) reported the utilisation of Eversa® Transform 2.0 to produce fatty ester for biolubricants. This study prepared several SFAEs from CPO and palm oil hydrolysates using Eversa® Transform 2.0 lipase as the catalyst. Commercially available pure palmitic acid, stearic acid and oleic acid were also employed to prepare SFAE as a comparison. Three types

of saccharides were used in this study, namely, D-mannitol (a sugar alcohol), D-mannose (an aldose) and D-fructose (a ketose), to better understand the relationship between the structural moiety, chemical reactivity and bioactivity. The feasibility of SFAE application in the health and cosmetics sector was explored by screening the synthesised SFAE for anticancer (cytotoxic) activity against MCF-7 cells and antimicrobial activity against *Cutibacterium acnes*. The MCF-7 cell line is a model for studying breast cancer, whereas *C. acnes* is associated with *Acne vulgaris*. Cancer is among the top ten causes of death worldwide, and breast cancer is among the top five deadly cancers (Sung et al., 2021).

MATERIALS AND METHODS

Fifteen SFAEs were included in this study. Fatty acids were obtained by the hydrolysis of CPO and palm oil. The palm oil was purchased from a local supermarket. Commercially pure ($\geq 98.0\%$) palmitic acid (Merck, Germany), stearic acid (Pudak Scientific, Indonesia) and oleic acid were used in this study. D-Mannitol and D-fructose were obtained from Sigma-Aldrich (USA) and D-mannose from HiMedia (India). The commercial lipase Eversa® Transform 2.0 from Novozyme (Denmark) was used as the catalyst in this study. Thin layer chromatography (TLC) analysis was conducted on TLC aluminium sheets coated with silica gel 60 F₂₅₄ using 2', 7'-dichlorofluorescein (Sigma-Aldrich) as the staining agent. All chemicals were used without further purification.

Hydrolysis of CPO and Palm Oil

The CPO and palm oil were hydrolysed separately under alkaline conditions. CPO and palm oil were separately mixed with a 1M NaOH solution in alcohol. The reaction mixture was refluxed for 1 hr and monitored using TLC. A solution of 1M HCl was added to the reaction mixture to neutralise the product. The product was extracted twice using *n*-hexane, dried using anhydrous Na₂SO₄ and concentrated to obtain CPO and palm oil hydrolysates. The fatty acid composition of CPO and palm oil was analysed using GC-MS. The acid and iodine numbers were also measured using the standard titrimetric method.

Synthesis of SFAE

Prior to the reaction, the reactants were dissolved in a suitable solvent. Fatty acids were dissolved separately in *n*-hexane, whereas the saccharides were dissolved in a small amount of water. Fatty acids were obtained by the hydrolysis of CPO, hydrolysis of palm oil and commercially available

fatty acids. Saccharides solution was added to the fatty acid solution followed by 5% weight-in-weight (w/w) lipase Eversa® Transform 2.0. The mole ratio of saccharide to fatty acid was 1:6. The mixture was incubated at 40°C for 48 hr and stirred at 500 rpm. The reaction mixture was heated at 100°C to deactivate the enzyme and stop the reaction. The product was separated from the reaction mixture by centrifugation. The product was washed with water to separate the enzyme and hexane to remove unreacted fatty acid. The amount of unreacted fatty acid was measured by acid-base titration using NaOH solution in alcohol. Conversion rate and degree of esterification were calculated by using the Equation (1) and (2):

$$\text{Conversion rate} = \frac{\text{Mole of initial fatty acid} - \text{Mole of unreacted fatty acid}}{\text{mole of initial fatty acid}} \times 100\% \quad (1)$$

$$\text{Degree of esterification} = \frac{\text{Mole of initial fatty acid} - \text{Mole of unreacted fatty acid}}{\text{Mole of saccharide}} \quad (2)$$

All SFAE was analysed by using Fourier transform infrared spectroscopy (FTIR) and only a representative of SFAE was analysed by using nuclear magnetic resonance (NMR) to confirm the formation of SFAE. FTIR spectra were analysed on potassium bromide (KBr) using a Shimadzu 2450 Prestige (Shimadzu Corp., Japan). ¹H-NMR and ¹³C-NMR spectra were recorded in CD₃OD (Merck, Germany) on a Bruker Avance Neo 500 (500 MHz) spectrophotometer (Bruker Biospin, Germany).

Emulsification Properties

The emulsification properties of SFAE were evaluated by adding 50 mg of SFAE to a water-oil mixture. A water-in-oil (w/o) emulsion was prepared by mixing 0.3 mL of water and 1 mL of oil, while a mixture of 0.3 mL of oil and 1 mL of water was used as a model for an oil-in-water (o/w) emulsion. The mixture containing SFAE was vortexed. The stability of the emulsions was observed for 24 hr. Phase separation was used as the parameter for emulsion stability. A light microscope and water-soluble dye (methylene blue) were used to facilitate observation of the emulsion. The emulsification properties of the starting material (saccharides and fatty acids) were also observed. A mixture of water and oil without a sample was used as the negative control.

Cytotoxic Activity Against MCF-7

MCF-7 cell suspension was added to 96-well plates at 5,000 cells/100 µL of medium. Dulbecco's Modified Eagle Medium (DMEM) supplemented

with 10% fetal bovine serum was used as the medium. The medium was enriched with 100 U/mL of penicillin and 100 µg/mL of streptomycin to prevent microbial contamination. The cells were incubated for 24 hr at 37°C before the addition of a series concentration of SFAE samples in dimethyl sulfoxide (DMSO). The final concentration of DMSO in the medium was 1% volume-in-volume (v/v). Therefore, 1% DMSO was used as the negative control. After 48 hr of incubation, 10 µL 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) (5 mg/mL) was added to each well. The plates were then incubated for another 4 hr. Ethanol was added to dissolve the formazan, and the sample absorption was read at 595 nm. The experiment was conducted in triplicate.

Antimicrobial Activity Against *C. acnes*

Antibacterial activity against *C. acnes* was determined using the broth dilution method in reinforced clostridial medium (RCM). A 250 mL *C. acnes* suspension was mixed with 5 mL RCM media and a 250 mL sample at room temperature. The DMSO was used as the SFAE solvent and the negative control. A 1,000 ppm salicylic acid was used as a positive control (the final concentration was 45 ppm). Antibacterial activity was determined by measuring optical density at 600 nm (OD₆₀₀). The experiment was conducted in duplicate.

RESULTS AND DISCUSSION

Fatty Acid Composition of CPO and Palm Oil

The fatty acid composition of palm oil and CPO was analysed using GC-MS (Table 1). CPO and palm oil showed similar fatty acid compositions. Oleic acid and palmitic acid were the major fatty acid constituents of CPO and palm oil, respectively. However, the oleic acid content of palm oil was slightly higher than that of CPO. Moreover, unsaturated medium-chain fatty acids (C8 and C10) were only detected in CPO but not in palm oil. Chompoo et al. (2019) reported that the degumming process during the conversion of CPO to palm oil increased the total amount of unsaturated fatty acids and reduced the total amount of saturated fatty acids.

Hydrolysates of CPO and Palm Oil

Hydrolysis of both CPO and palm oil was conducted under alkaline conditions. Acidic conditions were avoided to prevent changes in the C=C configuration of the unsaturated fatty acids in CPO and palm oil. The hydrolysate of CPO had a

lower R_f value ($R_f = 0.57$, hexane-ethyl acetate 3:1) on TLC than the starting material ($R_f = 0.86$, hexane-ethyl acetate 3:1), indicating the conversion of triglycerides to fatty acids. The TLC analysis of palm oil and its hydrolysate also showed a similar trend. The CPO hydrolysate was an orange solid, whereas the palm oil hydrolysate was a pale-yellow solid. Izuddin et al. (2022) stated that CPO is rich in β -carotene. Thus, the orange colour of CPO hydrolysate exhibited that the β -carotene could not be separated from the fatty acids. However, the presence of β -carotene in CPO hydrolysates did not interfere with the SFAE synthesis through the esterification reactions due to the absence of hydroxyl and carboxyl groups in its structure.

The formation of fatty acids from the hydrolysis of CPO and palm oil was also demonstrated by a significant increase in the acid number after hydrolysis (Table 2). However, the iodine numbers before and after hydrolysis were within similar ranges. It can be concluded that hydrolysis under alkaline conditions did not influence the unsaturated fatty acid content in CPO and palm oil. The electron-rich nature of the C=C bond is not reactive towards the nucleophilic property of the hydroxide anion from the alkaline hydrolysis reaction. The difference in chemical composition between CPO and palm oil hydrolysates affected the acid and iodine number values. CPO hydrolysate contained fatty acids and other nonpolar compounds (Goon et al., 2019) that cannot be separated during hydrolysis.

Synthesis of SFAE

The enzymatic reaction for preparing SFAE was conducted at 40°C according to Chang et al. (2021) and Farobie et al. (2021). Farobie et al. (2021)

stated that 40°C was the optimum temperature for enzymatic esterification using Eversa® Transform 2.0 lipase. The esterification reaction in this research was carried out in an organic phase with as little water as possible to promote the esterification and prevent the hydrolysis of the product. The water was used only to dissolve the sugar. The amount of water was calculated depending on the solubility of each saccharide. Therefore, molecular sieves were not used in the present study. Water adsorption by the molecular sieve separated the sugar from the reaction mixture. The esterification reaction catalysed by lipase occurs at the interface between the organic phase and water (Khan et al., 2017). Under hydrophobic conditions, the subdomain covering the lipase active site shifted, and the active site was accessible to the substrate. Hexane, which has a log P -value of 3.5, was chosen as the hydrophobic solvent in this study. Kumar et al. (2016) stated that solvents with log P -values above 2 did not interfere with the essential water in enzymes. Thus, hexane did not influence the active conformation of lipase.

The formation of SFAE was indicated by the appearance of a white layer between the water and hexane layers (Figure 1). The SFAE obtained in this research was sparingly insoluble in water or hexane at room temperature. The solubility test of sucrose fatty acid esters in water conducted by Nagai et al. (2017) showed that nine out of ten samples precipitated in water. Sucrose-lauric acid ester was the only sucrose ester that formed a transparent homogeneous solution (Nagai et al., 2017). The orange colour of the hexane layer in the SFAE of CPO hydrolysate (Figure 1a) denoted the presence of β -carotene. The β -carotene and SFAE of CPO hydrolysate can be separated easily at this stage.

TABLE 1. FATTY ACID COMPOSITION OF CRUDE PALM OIL (CPO) AND PALM OIL SAMPLES

Fatty acids	Percentage (%)	
	CPO	Palm oil
Caprylic acid (C8:0)	0.02	-
Capric acid (C10:0)	0.02	-
Lauric acid (C12:0)	0.18	0.21
Myristic acid (C14:0)	0.94	0.93
Palmitic acid (C16:0)	39.60	36.00
Stearic acid (C18:0)	4.59	4.08
Oleic acid (C18:1; 9c)	40.60	44.20

TABLE 2. ACID NUMBER AND IODINE NUMBER BEFORE AND AFTER HYDROLYSIS OF CRUDE PALM OIL (CPO) AND PALM OIL SAMPLES

Quality parameter	CPO		Palm oil	
	Before	After	Before	After
Acid number (mg NaOH/g sample)	28.6	204.8	5.3	183.1
Iodine number (mg I/g sample)	54.7	n.d	58.4	51.5




Figure 1. Formation of sugar fatty acid ester (SFAE) from (a) crude palm oil (CPO) and (b) palm oil hydrolysate.

Conversion Rate and Degree of Esterification

The esterification between carboxylic acids and alcohols is a reversible reaction. This study used an excessive amount of fatty acids to shift the equilibrium towards the formation of SFAE. The amount of reacted fatty acids was used to calculate the conversion rate and the degree of esterification (Table 3). The conversion rate and degree of esterification vary for each SFAE. In general, the Eversa® Transform 2.0 lipase used in this study preferred oleic acid, followed by stearic acid and palmitic acid. Depending on the source, lipases exhibit selectivity toward certain fatty acids (Halldorsson et al., 2004).

TABLE 3. CONVERSION RATE AND DEGREE OF ESTERIFICATION DURING THE SYNTHESIS OF SFAE

Sample's name	Conversion rate (%)	Degree of esterification
SFAE palmitic acid-mannitol	11.70	0.8
SFAE palmitic acid-mannose	21.40	1.3
SFAE palmitic acid-fructose	27.30	1.6
SFAE stearic acid-mannitol	30.30	1.8
SFAE stearic acid-mannose	20.60	1.2
SFAE stearic acid-fructose	13.90	0.8
SFAE oleic acid-mannitol	39.73	2.4
SFAE oleic acid-mannose	19.11	1.1
SFAE oleic acid-fructose	20.67	1.2
SFAE CPO-mannitol	74.80	4.4
SFAE CPO-mannose	58.10	2.0
SFAE CPO-fructose	61.00	3.7
SFAE palm oil-mannitol	22.40	1.3
SFAE palm oil-mannose	34.30	1.4
SFAE palm oil-fructose	21.80	2.0

Note: SFAE - sugar fatty acid esters; CPO - crude palm oil.

Most of the SFAE obtained in this study had a low degree of esterification. Zheng et al. (2015) reported that the products of enzymatic esterification commonly had a low degree of substitution. Lipase enzymes are more selective towards primary hydroxyl groups because of their low steric hindrance (Zago et al., 2021). Based on their structures, D-mannitol and D-fructose have two primary hydroxyl groups, whereas D-mannose only has one. Therefore, the degree of esterification of D-mannitol and D-fructose SFAE should be higher than that of D-mannose. SFAE of CPO hydrolysate, SFAE of oleic acid and SFAE of stearic acid showed the highest degree of esterification in D-mannitol. The hydroxyl groups of D-mannitol were assumed to be more reactive than those of D-fructose because D-mannitol does not form furanose or pyranose structures in solution. Meanwhile, the SFAE of palm oil hydrolysate and the SFAE of palmitic acid exhibited different behaviours, with D-fructose possessing the highest degree of esterification. Although the fatty acid compositions of CPO hydrolysate and palm oil hydrolysate were similar, the SFAE of CPO hydrolysate displayed a higher degree of esterification than SFAE-palm oil. The non-fatty acid components of the CPO hydrolysate may promote lipase activation. Sterol, vitamin E, dolichol and polyphenol are components of CPO (Goon et al., 2019) that may be responsible for lipase activation. Study by Delome et al. (2011) proved that the presence of surfactant can increase the activity of lipase by changing the three-dimensional structure of the enzyme and improving the accessibility of the substrate to the active site.

FTIR and NMR Spectra of SFAE

The formation of SFAE was confirmed by using FTIR and NMR spectra. The representative data of the FTIR spectra can be seen in Figure 2. The FTIR spectrum of the fatty acid (palmitic acid) showed a typical very broad hydroxyl peak at $3,400\text{--}2,500\text{ cm}^{-1}$ and a sharp carbonyl peak absorption of carboxylic acid at $1,703\text{ cm}^{-1}$. On the other hand, the hydroxyl group of the D-fructose and D-mannitol appeared as a broad peak at $3,400\text{--}3,000\text{ cm}^{-1}$. The absence of a very broad hydroxyl peak of the carboxylic group in the SFAE indicated the formation of an ester group between fatty acid (palmitic acid) and saccharides. Moreover, the shifting of the carbonyl peak to a higher frequency ($1,734\text{ cm}^{-1}$) also proved the presence of an ester group of SFAE.

The ^{13}C -NMR spectrum (Figure 3a) of SFAE palmitic acid-fructose exhibited the presence of carbonyl groups of esters at 175 and 176 ppm. The formation of an ester between palmitic acid and fructose can also be seen from the ^1H -NMR spectrum (Figure 3b). Based on the peak integral

value, it can be concluded that the degree of esterification of SFAE palmitic acid-fructose was more than one. The number of protons from the palmitoyl was about twice the number of protons from the D-fructose. However, the peak of fructose's acidic protons (the hydroxyl groups) cannot be seen clearly because the acidic protons are able to exchange with deuterated methanol as the analysis solvent. This data was suitable with the degree of esterification of SFAE palmitic acid-fructose.

Emulsification Properties of SFAE

The surfactant properties of the starting materials (fatty acids and saccharides) and

SFAE obtained in this study were examined by emulsification test in w/o and o/w systems. Emulsion stability was determined based on visual observation. The saccharides could not stabilise both w/o and o/w mixture. The o/w mixture with the addition of fatty acids gave a milky system that underwent phase separation after 15 min. The milky system consisting of a w/o system and fatty acids was stable for up to 30 min. Fameau et al. (2012) described that the fatty acids can reduce the surface tension.

The addition of SFAE prolonged the phase separation of the w/o and o/w system. All SFAE were more suitable for w/o emulsions than o/w emulsion systems. The phase separation was not




Figure 2. Fourier transform infrared spectroscopy (FTIR) spectra of (a) sugar fatty acid ester (SFAE) palmitic acid-fructose and (b) SFAE palmitic acid-mannitol compared to its starting material.




Figure 3. (a) ¹³C-NMR and (b) ¹H-NMR spectrum of sugar fatty acid ester (SFAE) palmitic acid-fructose.

observed on the w/o emulsion system containing SFAE after 24 hr. On the other hand, phase separation of the o/w system was noticed after 4 hr. SFAE possesses emulsifying properties because of the polar residues from sugar and nonpolar residues from fatty acids (Teng et al., 2021). Representative data are presented in Figure 4a and 4b. Figure 4c shows the microscopic observation of the w/o emulsion. Blue indicates the water phase.

Cytotoxic Activity Against MCF-7

The cytotoxic activity of SFAE was determined using an MTT assay based on mitochondrial enzyme activity in viable MCF-7 breast cancer cells (Perinelli et al., 2016). The cytotoxic activity of SFAE varied, with IC_{50} values between 29 and 200 ppm (Table 4). The SFAE of saturated fatty acids (palmitic and stearic acid) was more cytotoxic than the SFAE of oleic acid. These data are consistent with our previous research on the cytotoxicity of lipoamide derivatives. Lipoamides from saturated fatty acids exhibited the lowest IC_{50} value compared to lipoamides from unsaturated fatty acids (Wukirsari et al., 2024). The presence of the C=C bond is disadvantageous for cytotoxicity.

The cytotoxicity SFAE of D-mannose was lower than the SFAE of D-fructose and D-mannitol. The hydrophilic part SFAE of D-mannose (a pyranose ring) is bigger than D-fructose (a furanose ring) and D-mannitol (a linear chain). Among the SFAEs of D-fructose and D-mannitol synthesised from the same fatty acids, in general, the cytotoxic activity was inversely proportional to the degree of esterification. The SFAE of CPO hydrolysate with a higher degree of esterification also possessed lower cytotoxicity than the SFAE of palm oil hydrolysate. It seemed that the large size of SFAEs was unfavourable for SFAE-cell membrane interaction. According to Bunchongprasert and Shao (2020), the bulky size of the hydrophilic parts of fatty acid ester increased the intermolecular interactions of micelles and limited the SFAE interactions with the phospholipid bilayer

membrane of cancer cells. The disruption of cell membrane integrity is important for the cytotoxic activity of SFAE (Bunchongprasert & Shao, 2020). Okabe et al. (1999) stated that SFAE inhibited cancer cell proliferation by suppressing the secretion of TNF- α .

TABLE 4. CYTOTOXIC ACTIVITY OS SUGAR-FATTY ACID ESTERS (SFAE) AGAINST MCF-7 CELL LINE

Sample's name	IC_{50} (ppm)
SFAE palmitic acid-mannitol	29.5 ± 3.6
SFAE palmitic acid-mannose	95.9 ± 5.4
SFAE palmitic acid-fructose	78.4 ± 6.0
SFAE stearic acid-mannitol	54.3 ± 2.2
SFAE stearic acid-mannose	104.4 ± 7.2
SFAE stearic acid-fructose	29.7 ± 2.0
SFAE oleic acid-mannitol	127.4 ± 3.6
SFAE oleic acid-mannose	>200
SFAE oleic acid-fructose	125.2 ± 1.1
SFAE CPO-mannitol	131.0 ± 28.0
SFAE CPO-mannose	107.0 ± 11.0
SFAE CPO-fructose	66.0 ± 5.5
SFAE palm oil-mannitol	65.5 ± 1.0
SFAE palm oil-mannose	102.8 ± 3.0
SFAE palm oil-fructose	65.5 ± 1.0

Note: SFAE - sugar fatty acid esters; CPO - crude palm oil.

Antimicrobial Activity Against *C. acnes*

Using the broth dilution method, SFAE applications in the cosmetic sector were investigated by determining their antibacterial activity against the acne-causing bacterium *C. acnes*. The assay was conducted at a 0.9 mg/mL concentration for all samples. Bacterial growth was directly proportional to the optical density value. Therefore, samples with lower optical densities displayed higher antimicrobial activity. CPO hydrolysate, palm oil hydrolysate and SFAE showed antibacterial activity against *C. acnes* (Table 5). The sugar moiety of SFAE




Figure 4. Emulsification of sugar fatty acid ester (SFAE) properties in the (a) w/o and (b) o/w emulsion systems. (c) Observation of the w/o emulsion under a microscope; the blue colour indicates water dyed with methylene blue.

exhibited variable effects on antibacterial activity against *C. acnes*. Based on its cell wall composition, *C. acnes* is classified as a Gram-positive bacteria (Mayslich et al., 2021). Zhao et al. (2015) mentioned that Gram-positive bacteria were more sensitive towards SFAE than Gram-negative bacteria. Many researchers have also reported the antibacterial activity of fatty acids against *Propionibacterium acnes* (*C. acnes*) (Casillas-Vargas et al., 2021; Desbois & Lawlor, 2013; Kim et al., 2021). Several mechanisms for the antibacterial activity of fatty acids have been reported. Unsaturated fatty acids have been reported to prevent bacterial growth by interfering with DNA replication, cell wall formation and protein synthesis (Desbois & Lawlor, 2013). The disruption of bacterial cell membrane permeability by SFAE has been proposed by Zhao et al. (2015). Zhao et al. (2015) observed the leakage of bacterial cell components such as protein and reducing sugars after being treated with SFAE. Disruption of the microbe membrane is associated with the emulsification properties of SFAE (Falk, 2019; Pirog et al., 2019). The emulsification properties are influenced by the degree of esterification of SFAE and its HLB value (Chen et al., 2023). Therefore, the CPO and palm oil hydrolysate and also SFAE are promising candidate as active ingredient in anti-acne agents. The addition of SFAE as an ingredient in anti-acne products will provide dual benefits, i.e., as an antibacterial and emulsifier agent. The strong antimicrobial activity of SFAE was also stated by Yang et al. (2024).

TABLE 5. ANTIMICROBIAL ACTIVITY AGAINST *Cutibacterium acnes*

Sample's name	Inhibition (%)
CPO hydrolysate	86 ± 17
SFAE CPO-mannitol	100 ± 0
SFAE CPO-mannose	23
SFAE CPO-fructose	35 ± 12
Palm oil hydrolysate	100 ± 0
SFAE palm oil-mannitol	35 ± 9
SFAE palm oil-mannose	39 ± 9
SFAE palm oil-fructose	81 ± 6
Salicylic acid (45 mg/mL)	25 ± 1.7

Note: SFAE - sugar fatty acid esters; CPO - crude palm oil.

CONCLUSION

In this study, SFAE were synthesised through enzymatic esterification using the lipase enzyme Eversa® Transform 2.0. The potent cytotoxic effect was attributed to the saturated fatty constituents (acyl group) of the SFAE and SFAE having a low degree of esterification. SFAE of palmitic acid-mannitol and SFAE of stearic acid-fructose showed

the highest cytotoxic activity against MCF-7 breast cancer cells, with an IC_{50} value of 29 ppm. The SFAE of palm oil hydrolysate with a lower degree of esterification exhibited stronger cytotoxic activity than the SFAE of CPO hydrolysate. In contrast, SFAE from CPO hydrolysate and D-mannitol showed the strongest antibacterial activity achieving 100% inhibition against *C. acnes* at 0.9 mg/mL.

ACKNOWLEDGEMENT

This study was financially supported by the Research Grant Faculty of Mathematics and Natural Sciences, Universitas Indonesia (PKS-034/UN2.F3.D/PPM.00.02/2023). We would also like to thank AERG 23 (Alisya Rahmah, Angelia Sabrina Lexthariana, Novella Dhindha Pangestika, Olan Maulana and Zahrotun Nisa) for the support.

REFERENCES

- Bernal, C., Poveda-Jaramillo, J. C., & Mesa, M. (2018). Raising the enzymatic performance of lipase and protease in the synthesis of sugar fatty acid esters, by combined ionic exchange-hydrophobic immobilisation process on aminopropyl silica support. *Chemical Engineering Journal*, 334, 760–767. <https://doi.org/10.1016/j.cej.2017.10.082>
- Bunchongprasert, K., & Shao, J. (2020). Effect of fatty acid ester structure on cytotoxicity of self-emulsified nanoemulsion and transport of nanoemulsion droplets. *Colloids and Surfaces B: Biointerfaces*, 194, 111220. <https://doi.org/10.1016/j.colsurfb.2020.111220>
- Casillas-Vargas, G., Ocasio-Malavé, C., Medina, S., Morales-Guzmán, C., Del Valle, R. G., Carballeira, N. M., & Sanabria-Ríos, D. J. (2021). Antibacterial fatty acids: An update of possible mechanisms of action and implications in the development of the next-generation of antibacterial agents. *Progress in Lipid Research*, 82, 101093. <https://doi.org/10.1016/j.plipres.2021.101093>
- Chang, M. Y., Chan, E., & Song, C. P. (2021). Biodiesel production catalysed by low-cost liquid enzyme Eversa® Transform 2.0: Effect of free fatty acid content on lipase methanol tolerance and kinetic model. *Fuel*, 283, 119266. <https://doi.org/10.1016/j.fuel.2020.119266>
- Chen, J-Q., Hai, Y-W., Qing, C., Liang, M-Y., Banwell, M. G., & Lan, P. (2023). The synthesis of certain fatty acid ester derivatives of trehalose and an

- investigation of their emulsifying properties and bioactivities. *LWT – Food Science and Technology*, 187, 115369. <https://doi.org/10.1016/j.lwt.2023.115369>
- Chiewpattanakul, P., Phonnok, S., Durand, A., Marie, E., & Thanomsub, B. W. (2010). Bioproduction and anticancer activity of biosurfactant produced by the dematiaceous fungus *Exophiala dermatitidis* SK80. *Journal of Microbiology and Biotechnology*, 20(12), 1664–1671.
- Chompoo, M., Damrongwattanakool, N., & Raviyan, P. (2019). Effect of chemical degumming process on physicochemical properties of red palm oil. *Songklanakarin Journal of Science & Technology*, 41(3), 513–521.
- Delome, V., Dhouib, R., Canaan, S., Fotiadu, F., Carrière, F., & Cavalier, J.-F. (2011). Effect of surfactant on lipase structure, activity, and inhibition. *Pharmaceutical Research*, 28, 1831–1842. <https://doi.org/10.1007/s11095-010-0362-9>
- Desbois, A. P., & Lawlor, K. C. (2013). Antibacterial activity of long-chain polyunsaturated fatty acids against *Propionibacterium acnes* and *Staphylococcus aureus*. *Marine Drugs*, 11(11), 4544–4557. <https://doi.org/10.3390/md11114544>
- El-Baz, H. A., Elazzazy, A. M., Saleh, T. S., Dourou, M., Mahyoub, J. A., Baeshen, M. N., Madian, H. R., & Aggelis, G. (2021). Enzymatic synthesis of glucose fatty acid esters using SCOs as acyl group-donors and their biological activities. *Applied Sciences*, 11(6), 2700. <https://doi.org/10.3390/app11062700>
- Elmowafy, E., El-Derany, M. O., Biondo, F., Tiboni, M., Casettari, L., & Soliman, M. E. (2020). Quercetin loaded monolaurate sugar esters-based niosomes: Sustained release and mutual antioxidant-hepatoprotective interplay. *Pharmaceutics*, 12(2), 143. <https://doi.org/10.3390/pharmaceutics12020143>
- Falk, N. A. (2019). Surfactants as antimicrobials: A brief overview of microbial interfacial chemistry and surfactant antimicrobial activity. *Journal of Surfactants and Detergents*, 22(5), 1119–1127. <https://doi.org/10.1002/jsde.12293>
- Fameau, A.-L., Ventureira, J., Novales, B., & Douliez, J.-P. (2012). Foaming and emulsifying properties of fatty acids neutralized by tetrabutylammonium hydroxide. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 403, 87–95. <https://doi.org/10.1016/j.colsurfa.2012.03.059>
- Farobie, O., Jannah, Q. R., & Hartulistiyoso, E. (2021). Biodiesel production from crude palm oil under different free fatty acid content using Eversa® Transform 2.0 enzyme. *International Journal of Renewable Energy Research*, 11(4), 1590–1596. <https://doi.org/10.20508/ijrer.v11i4.12338.g8366>
- Ferdjani, S., & Selaimia, O. (2024). Synthesis of sugar esters using maize seed lipase as a biocatalyst. *Hungarian Journal of Industry and Chemistry*, 52(1), 9–14. <https://doi.org/10.33927/hjic-2024-02>
- Goon, D. E., Kadir, S. H. S. A., Latip, N. A., Rahim, S. A., & Mazlan, M. (2019). Palm oil in lipid-based formulations and drug delivery systems. *Biomolecules*, 9(2), 64. <https://doi.org/10.3390/biom9020064>
- Halldorsson, A., Kristinsson, B., & Haraldsson, G. G. (2004). Lipase selectivity toward fatty acids commonly found in fish oil. *European Journal of Lipid Science and Technology*, 106(2), 79–87. <https://doi.org/10.1002/ejlt.200300867>
- Hollenbach, R., Ochsenreither, K., & Syltatk, C. (2020). Enzymatic synthesis of glucose monodecanoate in a hydrophobic deep eutectic solvent. *International Journal of Molecular Sciences*, 21(12), 1–11. <https://doi.org/10.3390/ijms21124342>
- Izuddin, W. I., Loh, T. C., Akit, H., Nayan, N., Noor, A. M., & Foo, H. L. (2022). Influence of dietary palm oils, palm kernel oil and soybean oil in laying hens on production performance, egg quality, serum biochemicals and hepatic expression of beta-carotene, retinol and alpha-tocopherol genes. *Animals*, 12(22), 3156. <https://doi.org/10.3390/ani12223156>
- Khan, F. I., Lan, D., Durrani, R., Huan, W., Zhao, Z., & Wang, Y. (2017). The lid domain in lipases: Structural and functional determinant of enzymatic properties. *Frontiers in Bioengineering and Biotechnology*, 5, 16. <https://doi.org/10.3389/fbioe.2017.00016>
- Kim, Y. G., Lee, J. H., & Lee, J. (2021). Antibiofilm activities of fatty acids including myristoleic acid against *Cutibacterium acnes* via reduced cell hydrophobicity. *Phytomedicine*, 91, 153710. <https://doi.org/10.1016/j.phymed.2021.153710>
- Kumar, A., Dhar, K., Kanwar, S. S., & Arora, P. K. (2016). Lipase catalysis in organic solvents: Advantages and applications. *Biological Procedures Online*, 18, 2. <https://doi.org/10.1186/s12575-016-0033-2>

- Matin, M. M., Bhuiyan, M. M. H., Kibria, S. M., & Hasan, M. S. (2022). Synthesis, pass predication of antimicrobial activity and pharmacokinetic properties of hexanoyl galactopyranosides and experimental evaluation of their action against four human pathogenic bacteria and four fungal strains. *Pharmaceutical Chemistry Journal*, 56, 627–637. <https://doi.org/10.1007/s11094-022-02687-y>
- Matin, M. M., Hasan, M. S., Uzzaman, M., Bhuiyan, M. M. H., Kibria, S. M., Hossain, M. E., & Roshid, M. H. O. (2020). Synthesis, spectroscopic characterisation, molecular docking, and ADMET studies of mannopyranoside esters as antimicrobial agents. *Journal of Molecular Structure*, 1222, 128821. <https://doi.org/10.1016/j.molstruc.2020.128821>
- Mayslich, C., Grange, P. A., & Dupin, N. (2021). *Cutibacterium acnes* as an opportunistic pathogen: An update of its virulence-associated factors. *Microorganisms*, 9(2), 303. <https://doi.org/10.3390/microorganisms9020303>
- Monteiro, R. R. C., de Mela Neta, M. M. F., Rocha, W. S., Soares, J. B., de Luna, F. M. T., Fernandez-Lafuente, R., & Vieira, R. S. (2024). Optimizing the enzymatic production of biolubricants by the Taguchi method: Esterification of the free fatty acids from castor oil with 2-ethyl-1-hexanol catalyzed by Eversa Transform 2.0. *Enzyme and Microbial Technology*, 175, 110409. <https://doi.org/10.1016/j.enzmictec.2024.110409>
- Nagai, Y., Kawano, S., Motoda, K., Tomida, M., Tatebe, C., Sato, K., & Akiyama, H. (2017). Solubility testing of sucrose esters of fatty acids in international food additive specifications. *Biological and Pharmaceutical Bulletin*, 40(3), 284–289. <https://doi.org/10.1248/bpb.b16-00738>
- Nieto, S., Villa, R., Donaire, A., & Lozano, P. (2021). Ultrasound-assisted enzymatic synthesis of xylitol fatty acid esters in solvent-free conditions. *Ultrasonics Sonochemistry*, 75, 105606. <https://doi.org/10.1016/j.ultsonch.2021.105606>
- Okabe, S., Suganuma, M., Tada, Y., Ochiai, Y., Sueoka, E., Kohya, H., Shibata, A., Takahashi, M., Mizutani, M., Matsuzaki, T., & Fujiki, H. (1999). Disaccharide esters screened for inhibition of tumor necrosis factor- α release are new anti-cancer agents. *Japanese Journal of Cancer Research*, 90(6), 669–676. <https://doi.org/10.1111/j.1349-7006.1999.tb00799.x>
- Pauly, G., & Grisoni, P. (2005). *Use of sugar esters in cosmetic and/or pharmaceutical preparations*. (U.S. Patent No. 2005/0220726 A1). U.S. Patent Application Publication.
- Perinelli, D. R., Casettari, L., Cespi, M., Fini, F., Man, D. K. W., Giorgioni, G., Canala, S., Lam, J. K. W., Bonacucina, G., & Palmieri, G. F. (2016). Chemical-physical properties and cytotoxicity of *N*-decanoyl amino acid-based surfactants: Effect of polar heads. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 492, 38–46. <https://doi.org/10.1016/j.colsurfa.2015.12.009>
- Pirog, T. P., Lutsay, D. A., Kliuchka, L. V., & Beregova, K. A. (2019). Antimicrobial activity of surfactants of microbial origin. *Biotechnologia Acta*, 12(1), 30–57. <https://doi.org/10.15407/biotech.12.01.039>
- Qi, Y., Chen, M., Jin, T., Chong, W., Zhang, Z., Nian, B., & Hu, Y. (2024). Computer-aided engineering of lipase solvent tolerance enhanced their applications in sugar esters industry: State of the art. *Trends in Food Science and Technology*, 144, 104323. <https://doi.org/10.1016/j.tifs.2023.104323>
- Sardar, N. R., Akabari, S. H., Bhatt, H. G., & Modi, R. B. (2024). Role of sucrose ester for enhancing the food quality and safety: A review. *Asian Journal of Food Research and Nutrition* 3(3), 425–430. <https://journalajfrn.com/index.php/AJFRN/article/view/144>
- Shin, D. W., Mai, N. L., Bae, S. W., & Koo, Y. M. (2019). Enhanced lipase-catalysed synthesis of sugar fatty acid esters using supersaturated sugar solution in ionic liquids. *Enzyme and Microbial Technology*, 126, 18–23. <https://doi.org/10.1016/j.enzmictec.2019.03.004>
- Snoch, W., Wnuk, D., Witko, T., Staroń, J., Bojarski, A. J., Jarek, E., Plou, F. J., & Guzik, M. (2021). In search of effective anticancer agents-novel sugar esters based on polyhydroxyalkanoate monomers. *International Journal of Molecular Sciences*, 22(13), 7238. <https://doi.org/10.3390/ijms22137238>
- Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., & Bray, F. (2021). Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians*, 71(3), 209–249. <https://doi.org/10.3322/caac.21660>
- Teng, Y., Stewart, S. G., Hai, Y. W., Li, X., Banwell, M. G., & Lan, P. (2021). Sucrose fatty acid esters:

- Synthesis, emulsifying capacities, biological activities and structure-property profiles. *Critical Reviews in Food Science and Nutrition*, 61(19), 3297–3317. <https://doi.org/10.1080/10408398.2020.1798346>
- U.S. Department of Agriculture. (2024, October 7). *Palm oil production: Top producing countries*. <https://fas.usda.gov/data/production/commodity/4243000>
- Vitiello, L., Lixi, F., Coco, G., & Giannaccare, G. (2024). Ocular surface side effect of novel anticancer drugs. *Cancer*, 16(2), 344. <https://doi.org/10.3390/cancers16020344>
- Walvekar, S., Yasaswi, S., Shetty, K., & Yadav, K. S. (2022). Applications of surfactin and other biosurfactants in anticancer activity. In *Green sustainable process for chemical and environmental engineering and science* (pp. 223–234). Elsevier. <https://doi.org/10.1016/B978-0-323-85146-6.00024-3>
- Wang, Y., Wang, Z., Sun, Y., Zhu, M., Jiang, Y., Bai, H., Yang, B., & Kuang, H. (2024). Isovaleryl sucrose esters from *Atractylodes japonica* and their cytotoxic activity. *Molecules*, 29(13), 3069. <https://doi.org/10.3390/molecules29133069>
- Wiśniewska, J., Klasik-Ciszewska, S., & Duda-Grychtoł, K. (2023). Salicylic acid and its use in cosmetology. *Aesthetic Cosmetology and Medicine*, 12(3), 91–95. <https://doi.org/10.52336/acm.2023.011>
- Wukirsari, T., Nadiroh, N., Kembara, T. F., Gunawan, M. I. A., Priosoeryanto, B. P., Handayani, S., & Hudiyono, S. (2024). The effect of fatty alkanolamide structure on cytotoxic and antimicrobial activity. *Egyptian Journal of Chemistry*, 67(9), 289–295. <https://doi.org/10.21608/ejchem.2023.212786.8007>
- Yang, J., Huang, D., Wang, X., Liu, Q., & Guo, R. (2024). Research progress of eco-friendly sugar-based surfactants. *Academic Journal of Science and Technology*, 10(2), 70–77. <https://doi.org/10.54097/002vp898>
- Zago, E., Joly, N., Chaveriat, L., Lequart, V., & Martin, P. (2021). Enzymatic synthesis of amphiphilic carbohydrate esters: Influence of physicochemical and biochemical parameters. *Biotechnology Reports*, 30, e00631. <https://doi.org/10.1016/j.btre.2021.e00631>
- Zhao, L., Zhang, H., Hao, T., & Li, S. (2015). *In vitro* antibacterial activities and mechanism of sugar fatty acid esters against five food-related bacteria. *Food Chemistry*, 187, 370–377. <https://doi.org/10.1016/j.foodchem.2015.04.108>
- Zheng, Y., Zheng, M., Ma, Z., Xin, B., Guo, R., & Xu, X. (2015). Sugar fatty acid esters. *Polar Lipids*, 215–243. <https://doi.org/10.1016/b978-1-63067-044-3.50012-1>